

26-0-3 MESA



GUARANTEED ANALYSIS	GUARANTEED MINIMUM ANALYSIS
Total Nitrogen (N) 26%	Total Nitrogen (N)
4.8% Ammoniacal Nitrogen	4.8% Ammoniacal Nitrogen
3.6% Water Insoluble Nitrogen*	3.6% Water Insoluble Nitrogen*
12.4% Urea Nitrogen	12.4% Urea Nitrogen
5.2% Other Water Soluble Nitrogen*	5.2% Other Water Soluble Nitrogen*
Soluble Potash (K ₂ O)	
Sulfur (S)	Sultur (S)
5.7% Combined Sulfur (S)	5.7% Combined Sulfur (S)
Iron (Fe)	1107/0
0.18% Water Insoluble Iron (Fe)	0.18% Water Insoluble Iron (Fe)
Derived From: methylene ureas, ammonium sulfate, urea, biosolids, sulfate of potasl	Derived From: methylene ureas, ammonium sulfate, urea, biosolids, sulfate of potash.
*0.7% slowly available nitrogen from biosolids.	Chlorine (CI) maximum 1.5%
*8.1% slowly available nitrogen from methylene ureas.	31101110 (31) Haxillani 11370
, , , ,	*0.7% slowly available nitrogen from biosolids.
Chlorine (CI) not more than	*8.1% slowly available nitrogen from methylene ureas.
U.S.A.	Canada

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.5 lbs per 1,000 sq. ft. or 150 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 1.9 lbs per 1,000 sq. ft. or 83 lbs. per

SUGGESTED SPREADER SETTINGS			
Spreader	0.9#N /1,000 sq. ft.	0.5#N /1,000 sq. ft.	
Lebanon Turf	51/4	41/2	
Andersons AccuPr	о О	K	
Earthway Rotary	19	15	
Gandy	31	26	
Lely	6½ II	5 II	
Lesco (letter/nume	ric) K/23	i/17	
PennMulch HVO	0	L	
ProScape SS	0	L	
Spyker	51/4	41/2	
Vicon	30	24	
PennMulch HVO ProScape SS Spyker	O O 51/4	L L 4½	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 1.8 kg of product per 100 sq. meters.

To feed at the rate of 0.23 kg Nitrogen (N) Equivalent to 1.0 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS			
Spreader 0	.43 kg N/100 m ²	0.23 kg# N/100 m ²	
Lebanon Turf	51/4	41/2	
Andersons AccuPro	0	K	
Earthway Rotary	19	15	
Gandy	31	26	
Lely	6½ II	5 II	
Lesco (letter/numeric) K/23	i/17	
PennMulch HVO	0	L	
ProScape SS	0	L	
Spyker	51/4	41/2	
Vicon	30	24	
These settings were calibrated and field tested			

These settings were calibrated and field tested.

However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Hazard Statements:

May cause skin irritation. May be harmful if inhaled. May cause eye irritation.



Precautionary Statements for handling:

Wash hands and exposed skin thoroughly after handling.

Wear protective gloves, protective clothing, eye protection, and face protection.

If eye exposure occurs, rinse with water and seek medical attention.

Take off contaminated clothing and wash before reuse.

If skin irritation occurs seek medical advice

Store locked up in a well-ventilated place.

Keep the container tightly closed and out of reach of children.

Dispose of in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn. Do not apply to frozen ground.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

Do not exceed 2.0 lbs. total phosphorus annually

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the

states of Maryland, New Jersey, New York or Virginia except when
1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application

2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State

3) Re-establishing or repairing a turf area.

*In Maryland, the soil testing laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms. **In Pennsylvania:** Do not exceed 0.9 lbs of total nitrogen per 1,000 sq. ft. per application.



Manufactured by: Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042 800-233-0628 • (717) 273-1685 www.LebanonTurf.com