

LebanonTurf

Country Club[®] MD

Maximum Dispersion

- Utilizes a unique, highly dispersing granule technology
- Contains North Atlantic Ocean sea kelp
- Features METH-EX[®] slow release nitrogen
- SGN 80 is ideal for putting greens

FERTILIZER

11-7-21



GUARANTEED ANALYSIS

Total Nitrogen (N).....	11%
1.5% Ammoniacal Nitrogen	
2.0% Water Insoluble Nitrogen*	
3.9% Urea Nitrogen	
3.6% Other Water Soluble Nitrogen *	
Available Phosphate (P ₂ O ₅).....	7%
Soluble Potash (K ₂ O).....	21%
Magnesium (Mg).....	1.0%
1.0% Water Soluble Magnesium (Mg)	
Sulfur (S).....	8.4%
8.4% Combined Sulfur (S)	
Iron (Fe).....	2.0%
0.2% Water Soluble Iron (Fe)	
Manganese (Mn).....	1.0%
1.0% Water Soluble Manganese (Mn)	

Derived From: Ammonium Phosphate, Methylene Ureas, Urea, Kelp Meal, Sulfate of Potash, Sulfate of Potash Magnesia, Ferric Oxide, Ferrous Sulfate, Manganese Sulfate.
*5.6% Slowly Available Nitrogen from Methylene Ureas.

Chlorine (Cl) not more than..... 2.0%

ALSO CONTAINS NON PLANT FOOD INGREDIENTS
SOIL/PLANT AMENDING INGREDIENTS
Active Ingredients:

Humic Acids (derived from Leonardite).....	1.0%
Inert Ingredients (as soil amending).....	99.0%
99.0% Fertilizer Ingredients	

Country Club MD products contain humic acids, which may aid in nutrient availability.

U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N).....	11%
1.5% Ammoniacal Nitrogen	
2.0% Water Insoluble Nitrogen*	
3.9% Urea Nitrogen	
3.6% Other Water Soluble Nitrogen *	
Available Phosphate (P ₂ O ₅).....	7%
Soluble Potash (K ₂ O).....	21%
Magnesium (Mg).....	1.0%
1.0% Water Soluble Magnesium (Mg)	
Sulfur (S).....	8.4%
8.4% Combined Sulfur (S)	
Iron (Fe) (actual).....	2.0%
0.2% Water Soluble Iron (Fe)	
Manganese (Mn) (actual).....	1.0%
1.0% Water Soluble Manganese (Mn)	
Chlorine (Cl) (Maximum).....	2.0%

Derived From: Ammonium Phosphate, Methylene Ureas, Urea, Kelp Meal, Sulfate of Potash, Sulfate of Potash Magnesia, Ferric Oxide, Ferrous Sulfate, Manganese Sulfate.
*5.6% Slowly Available Nitrogen from Methylene Ureas.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS
SOIL/PLANT AMENDING INGREDIENTS
Active Ingredients:

Humic Acids (derived from Leonardite).....	1.0%
Inert Ingredients (as soil amending).....	99.0%
99.0% Fertilizer Ingredients	

Country Club MD products contain humic acids, which may aid in nutrient availability.

CANADA

NOTICE: This product contains the secondary micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.
Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:
To feed at the rate of 1 lb. Potash (K₂O) per 1,000 sq. ft., apply this product at 4.8 lbs per 1,000 sq. ft. or 209 lbs. per acre. To feed at the rate of ½ lb. Potash (K₂O) per 1,000 sq. ft., apply this product at 2.4 lbs per 1,000 sq. ft. or 105 lbs. per acre.

Spreader	1#K ₂ O/1,000 sq. ft.	0.5#K ₂ O/1,000 sq. ft.
Lebanon Turf	3	2
Andersons AccuPro	H	F
Earthway Rotary	13	11
Gandy	23	21
Lely	3½ II	2½ II
Lesco (letter/numeric)	G/16	E/15
PennMulch HVO	i	G
ProScape SS	i	G
Spyker	3	1½
Vicon	20	16

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application;

2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State;

or 3) Re-establishing or repairing a turf area.
*In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

Warning

Hazard Statements
May be harmful if inhaled repeatedly over prolonged periods.
Causes skin irritation.
Causes eye irritation.
May be harmful if inhaled repeatedly over prolonged periods.
May cause respiratory irritation.
Suspected of causing cancer by prolonged periods.
Harmful if swallowed.



Precautionary Statements
Keep container tightly closed and store in a well-ventilated place.
Avoid breathing dust.
Wash hands and exposed skin thoroughly after handling.
Use appropriate personal protective equipment as required to avoid breathing dust.
Do not eat, drink or smoke when using this product.
Use only outdoors or in a well-ventilated area.
Wear protective gloves/protective clothing/eye protection/face protection.
IF SWALLOWED: Rinse mouth. Immediately call a POISON CENTER or doctor/physician.
IF ON SKIN: Gently wash with plenty of soap and water. Seek medical attention if irritation occurs and persists.
If in eyes, rinse cautiously with water for several minutes. If eye irritation persists: Seek medical advice.
Call a poison center or doctor/physician if you feel unwell.
If inhaled and breathing difficulty occurs, remove victim to fresh air and keep at rest in a position comfortable for breathing.
Take off contaminated clothing and wash before reuse.
Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:
To feed at the rate of 0.48 kg Potash (K₂O) Equivalent to 2.3 kg of product per 100 sq. meters.
To feed at the rate of 0.24 kg Potash (K₂O) Equivalent to 1.2 kg of product per 100 sq. meters.

Spreader	0.48kg #K ₂ O/100 m ² .	0.24kg #K ₂ O/100 m ² .
Lebanon Turf	3	2
Andersons AccuPro	H	F
Earthway Rotary	13	11
Gandy	23	21
Lely	3½ II	2½ II
Lesco (letter/numeric)	G/16	E/15
PennMulch HVO	i	G
ProScape SS	i	G
Spyker	3	1½
Vicon	20	16

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

21-29078

C/17

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation
1600 E. Cumberland St • Lebanon, PA 17042
800-233-0628 • (717-273-1685)

40 Lbs. (18.14 kg) Net Weight

LebanonTurf



Country Club[®] MD

Maximum Dispersion

- Utilizes a unique, highly dispersing granule technology
- Contains North Atlantic Ocean sea kelp
- Features METH-EX[®] slow release nitrogen
- SGN 80 is ideal for putting greens

FERTILIZER

11-7-21



GUARANTEED ANALYSIS

Total Nitrogen (N)	11%
1.5% Ammoniacal Nitrogen	
2.0% Water Insoluble Nitrogen*	
3.9% Urea Nitrogen	
3.6% Other Water Soluble Nitrogen *	
Available Phosphate (P ₂ O ₅)	7%
Soluble Potash (K ₂ O)	21%
Magnesium (Mg).....	1.0%
1.0% Water Soluble Magnesium (Mg)	
Sulfur (S).....	8.4%
8.4% Combined Sulfur (S)	
Iron (Fe)	2.0%
0.2% Water Soluble Iron (Fe)	
Manganese (Mn).....	1.0%
1.0% Water Soluble Manganese (Mn)	

Derived From: Ammonium Phosphate, Methylene Ureas, Urea, Kelp Meal, Sulfate of Potash, Sulfate of Potash Magnesia, Ferric Oxide, Ferrous Sulfate, Manganese Sulfate.
*5.6% Slowly Available Nitrogen from Methylene Ureas.
Chlorine (Cl) not more than..... 2.0%

ALSO CONTAINS NON PLANT FOOD INGREDIENTS

SOIL/PLANT AMENDING INGREDIENTS

Active Ingredients:

Humic Acids (derived from Leonardite)	1.0%
Inert Ingredients (as soil amending)	99.0%
99.0% Fertilizer Ingredients	

U.S.A

Country Club MD products contain humic acids, which may aid in nutrient availability.

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	11%
1.5% Ammoniacal Nitrogen	
2.0% Water Insoluble Nitrogen*	
3.9% Urea Nitrogen	
3.6% Other Water Soluble Nitrogen *	
Available Phosphate (P ₂ O ₅)	7%
Soluble Potash (K ₂ O)	21%
Magnesium (Mg).....	1.0%
1.0% Water Soluble Magnesium (Mg)	
Sulfur (S).....	8.4%
8.4% Combined Sulfur (S)	
Iron (Fe) (actual).....	2.0%
0.2% Water Soluble Iron (Fe)	
Manganese (Mn) (actual)	1.0%
1.0% Water Soluble Manganese (Mn)	
Chlorine (Cl) (Maximum).....	2.0%

Derived From: Ammonium Phosphate, Methylene Ureas, Urea, Kelp Meal, Sulfate of Potash, Sulfate of Potash Magnesia, Ferric Oxide, Ferrous Sulfate, Manganese Sulfate.
*5.6% Slowly Available Nitrogen from Methylene Ureas.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS

SOIL/PLANT AMENDING INGREDIENTS

Active Ingredients:

Humic Acids (derived from Leonardite)	1.0%
Inert Ingredients (as soil amending)	99.0%
99.0% Fertilizer Ingredients	

CANADA

Country Club MD products contain humic acids, which may aid in nutrient availability.

NOTICE: This product contains the secondary micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 1 lb. Potash (K₂O) per 1,000 sq. ft., apply this product at 4.8 lbs per 1,000 sq. ft. or 209 lbs. per acre. To feed at the rate of ½ lb. Potash (K₂O) per 1,000 sq. ft., apply this product at 2.4 lbs per 1,000 sq. ft. or 105 lbs. per acre.

SUGGESTED SPREADER SETTINGS

Spreader	1#K ₂ O/1,000 sq. ft.	0.5#K ₂ O/1,000 sq. ft.
Lebanon Turf	3	2
Andersons AccuPro	H	F
Earthway Rotary	13	11
Gandy	23	21
Lely	3½ II	2½ II
Lesco (letter/numeric)	G/16	E/15
PennMulch HVO	i	G
ProScape SS	i	G
Spyker	3	1½
Vicon	20	16

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

- 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application;
 - 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State;
 - or 3) Re-establishing or repairing a turf area.
- *In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

Warning

Hazard Statements

May be harmful if inhaled repeatedly over prolonged periods.
Causes skin irritation.
Causes eye irritation.
May be harmful if inhaled repeatedly over prolonged periods.
May cause respiratory irritation.
Suspected of causing cancer by prolonged periods.
Harmful if swallowed.

Precautionary Statements

Keep container tightly closed and store in a well-ventilated place.
Avoid breathing dust.
Wash hands and exposed skin thoroughly after handling.
Use appropriate personal protective equipment as required to avoid breathing dust. Do not eat, drink or smoke when using this product.
Use only outdoors or in a well-ventilated area.
Wear protective gloves/protective clothing/eye protection/face protection.
IF SWALLOWED: Rinse mouth. Immediately call a POISON CENTER or doctor physician.
IF ON SKIN: Gently wash with plenty of soap and water. Seek medical attention if irritation occurs and persists.
If in eyes, rinse cautiously with water for several minutes. If eye irritation persists: Seek medical advice.
Call a poison center or doctor/physician if you feel unwell.
If inhaled and breathing difficulty occurs, remove victim to fresh air and keep at rest in a position comfortable for breathing.
Take of contaminated clothing and wash before reuse.
Dispose in accordance with all federal, state and local regulations.



Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.48 kg Potash (K₂O) Equivalent to 2.3 kg of product per 100 sq. meters.

To feed at the rate of 0.24 kg Potash (K₂O) Equivalent to 1.2 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

Spreader	0.48kg #K ₂ O/100 m ² .	0.24kg #K ₂ O/100 m ² .
Lebanon Turf	3	2
Andersons AccuPro	H	F
Earthway Rotary	13	11
Gandy	23	21
Lely	3½ II	2½ II
Lesco (letter/numeric)	G/16	E/15
PennMulch HVO	i	G
ProScape SS	i	G
Spyker	3	1½
Vicon	20	16

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.


Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>



F699

21-29078

C/17

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042

800-233-0628 • (717-273-1685)

40 Lbs. (18.14 kg) Net Weight