LebanonTurl

Country Club® Blended Fertilizer

19-0-19



GUARANTEED ANALYSIS

Total Nitrogen (N)	19%
3.6% Ammoniacal Nitrogen	
4.8% Water Insoluble Nitrogen*	
2.1% Urea Nitrogen	
8.5% Other Water Soluble Nitrogen *	
Soluble Potash (K ₂ 0)	19%
Sulfur (S)	13%
13.0% Combined Sulfur (S)	
Iron (Fe)	3.3%
0.3% Water Soluble Iron (Fe)	
Derived From: Methylene Ureas, Ammonium Sulfate, Ferric Oxide, Ferro	ous Sulfate.
*13.3% Slowly Available Nitrogen from Methylene Ureas.	
Chlorine (CI) not more than	2.0%
U	I.S.A

3.6% Ammoniacal Nitrogen		
4.8% Water Insoluble Nitrogen*		
2.1% Urea Nitrogen		
8.5% Other Water Soluble Nitrogen *		
Soluble Potash (K ₂ 0)	19%	
Sulfur (S)	13%	
13.0% Combined Sulfur (S)		
Iron (Fe) (actual)	3.3%	
0.3% Water Soluble Iron (Fe)		
Chlorine (CI) (maximum)		
Derived From: Methylene Ureas, Ammonium Sulfate, Ferric Oxide, Ferrous Sulfate. *13.3% Slowly Available Nitrogen from Methylene Ureas.		

CANADA

Notice: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after applications by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

WARNING

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.7 lbs per 1,000 sq. ft. or 204 lbs. per acre. To feed at the rate of $\frac{1}{2}$ lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.69 lbs per 1,000 sq. ft. or 113 lbs. per acre.

SUGGESTED SPREADER SETTINGS			
Spreader	0.9# N/1,000 sq. ft.	0.5 # N/1,000 sq. ft.	
Lebanon Turf	5	4	
Andersons Aco	cuPro N	J	
Earthway Rota	ry 17	14	
Gandy	29	25	
Lely	6 II	4½ II	
Lesco (letter/r	iumeric) J/21	H/16	
PennMulch HV	70 N	K	
ProScape SS	N	K	
Spyker .	5	4	
Vicon	28	23	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Hazard Statements:

Causes skin irritation.
May be harmful if inhaled.
Causes serious eye irritation.

Precautionary Statements for handling:

Wash hands and exposed skin thoroughly after handling.

Wear protective gloves/protective clothing/eye protection/face protection.

Take off contaminated clothing and wash before reuse.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. P351: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If skin irritation occurs: Get medical advice/ attention.

If eye irritation persists: Get medical advice/attention.

Keep out of reach of children

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 $\mbox{m}^2,$ applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 0.23 kg of product per 100 sq. meters.

To feed at the rate of 0.23 kg Nitrogen (N) Equivalent to 0.12 kg of product per 100 sq. meters.

	3 - 1 1		
SUGGESTED SPREADER SETTINGS			
Spreader	0.43 kg N /100 m ²	0.22 kg N /100 m ²	
Lebanon Turf	5	4	
Andersons Accı	iPro N	J	
Earthway Rotar	y 17	14	
Gandy	29	25	
Lely	6 II	4½ II	
Lesco (letter/nu	meric) J/21	H/16	
PennMulch HV0	N	K	
ProScape SS	N	K	
Spyker	5	4	
Vicon	28	23	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

