LebanonTurf

Country Club® Blended Fertilizer

17-0-17



GUARANTEED ANALYSIS

| Total Nitrogen (N) | 17% |
|---|------|
| 5.4% Water Insoluble Nitrogen* | |
| 2.2% Urea Nitrogen | |
| 9.4% Other Water Soluble Nitrogen * | |
| Soluble Potash (K ₂ 0) | |
| Sulfur (S) | 5.8% |
| 5.8% Combined Sulfur (S) | |
| Iron (Fe) | 0.5% |
| 0.005% Water Soluble Iron (Fe) | |
| | |
| Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate. | |
| | |
| *14.8% Slowly Available Nitrogen from Methylene Ureas. | |
| | |
| Chlorine (CI) not more than | 2.0% |

determined by the professional applicator based on local soils, regulations, and climate.

GUARANTEED MINIMUM ANALYSIS

| Total Nitrogen (N) | 17% |
|---|------|
| Soluble Potash (K ₂ 0) | 17% |
| Sulfur (S) | 5.8% |
| 5.8% Combined Sulfur (S) | |
| Iron (Fe) (actual) | 0.5% |
| 0.005% Water Soluble Iron (Fe) | |
| Chlorine (CI) (maximum) | 2.0% |
| Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate. | |

*14.8% Slowly Available Nitrogen from Methylene Ureas.

CANADA

Notice: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after applications by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 5.3 lbs per 1,000 sq. ft. or 230 lbs. per acre. To feed at the rate of ½ lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.9 lbs per 1,000 sq. ft. or 126 lbs. per acre.

| 11. 01 120 103. pei acie. | | |
|-----------------------------|----------------------|---------------------|
| SUGGESTED SPREADER SETTINGS | | |
| Spreader | 0.9# N/1,000 sq. ft. | 0.5 # N/1,000 sq. f |
| Lebanon Turf | 41/2 | 31/2 |
| Andersons Ac | cuPro K | Н |
| Earthway Rot | ary 16 | 14 |
| Gandy | 28 | 24 |
| Lely | 6 II | 4 II |
| Lesco (letter/ | numeric) i/17 | H/15 |
| PennMulch H | VO L | J |
| ProScape SS | L | J |
| Spyker | 41/2 | 31/2 |
| Vicon | 24 | 22 |
| | | |

These settings were calibrated and field tested. However, age and condition of spreader, speed of peration and evenness of terrain may require slightly different settings for desired coverage.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions



Hazard Statements Causes skin irritation.

Causes eye irritation. May be harmful if inhaled.

Suspected of causing cancer

Precautionary Statements for handling:

Do not handle until all safety precautions have been read and understood. Avoid breathing dust/fume/gas/mist/vapours/spray Wash hands and exposed skin thoroughly after handling

Wear protective gloves/protective clothing/eye protection/face protection. Use personal protective equipment as required. If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse.

IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Call POISON CENTER or doctor/physician if you feel unwell. If skin irritation occurs: Get medical advice/ attention.

If eye irritation persists: Get medical advice/attention. Store locked up. Dispose of contents/container properly as required by law for waste pesticides

If exposed or concerned, seek medical advice. Dispose in accordance with all federal, state and local regulations. Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.6 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates: To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 2.5 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N) Equivalent to 1.4 kg of product per 100 sq. meters.

| | <u> </u> | • | |
|-----------------------------|-------------------------------|-------------------------------|--|
| SUGGESTED SPREADER SETTINGS | | | |
| Spreader | 0.43 kg N /100 m ² | 0.22 kg N /100 m ² | |
| Lebanon Turf | 41/2 | 31/2 | |
| Andersons Acc | uPro K | Н | |
| Earthway Rotai | y 16 | 14 | |
| Gandy | 28 | 24 | |
| Lely | 6 II | 4 II | |
| Lesco (letter/ni | umeric) i/17 | H/15 | |
| PennMulch HV |) L | J | |
| ProScape SS | L | J | |
| Spyker | 41/2 | 31/2 | |
| | | | |

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

PennMulch® and ProScape®. Country Club and the Lebanon Diamond are registered trademarks of Lebanon Seaboard Corporation.



For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685



GUARANTEED ANALYSIS

| Total Nitrogen (N) | 17% |
|---|------|
| 9.4% Other Water Soluble Nitrogen * | |
| Soluble Potash (K ₂ O) | 17% |
| Sulfur (S) | 5.8% |
| 5.8% Combined Sulfur (S) | |
| Iron (Fe) | 0.5% |
| Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate. | |
| *14.8% Slowly Available Nitrogen from Methylene Ureas. | |
| Chlorine (CI) not more than | 2.0% |

| CHARA | NTEED | MINIMUM | ΔΝΔΙΥΟΙΟ |
|--------|-------|---------|------------|
| UUAIIA | | | AIIALI DID |

| Total Nitrogen (N) 5.4% Water Insoluble Nitrogen* | 17% |
|--|------|
| 2.2% Urea Nitrogen | |
| 9.4% Other Water Soluble Nitrogen * | |
| Soluble Potash (K ₂ 0) | |
| Sulfur (S) | 5.8% |
| 5.8% Combined Sulfur (S) | |
| Iron (Fe) (actual) | 0.5% |
| 0.005% Water Soluble Iron (Fe) | |
| Chlorine (CI) (maximum) | 2.0% |
| Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate |). |
| | |

*14.8% Slowly Available Nitrogen from Methylene Ureas.

CANADA

Notice: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after applications by sweeping or blowing. Do not wash off with water.

U.S.A

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 5.3 lbs per 1,000 sq. ft. or 230 lbs. per acre. To feed at the rate of $\frac{1}{2}$ lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.9 lbs per 1,000 sq. ft. or 126 lbs. per acre.

| SUGGESTED SPREADER SETTINGS | | | |
|-----------------------------|----------------------|-----------------------|--|
| Spreader | 0.9# N/1,000 sq. ft. | 0.5 # N/1,000 sq. ft. | |
| Lebanon Turf | 41/2 | 31/2 | |
| Andersons Acc | uPro K | H | |
| Earthway Rotar | y 16 | 14 | |
| Gandy | 28 | 24 | |
| Lely | 6 II | 4 II | |
| Lesco (letter/nu | ımeric) i/17 | H/15 | |
| PennMulch HV(|) L | J | |
| ProScape SS | L | J | |
| Spyker | 41/2 | 31/2 | |
| Vicon | 24 | 22 | |
| | | | |

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor. agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.



Hazard Statements

Causes skin irritation. Causes eye irritation. May be harmful if inhaled. Suspected of causing cancer.

Precautionary Statements for handling:

Do not handle until all safety precautions have been read and understood. Avoid breathing dust/fume/gas/mist/vapours/spray.

Wash hands and exposed skin thoroughly after handling.

Wear protective gloves/protective clothing/eye protection/face protection. Use personal protective equipment as required.

If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call POISON CENTER or doctor/physician if you feel unwell.

If skin irritation occurs: Get medical advice/ attention.

If eye irritation persists: Get medical advice/attention.

Store locked up.

Dispose of contents/container properly as required by law for waste pesticides. If exposed or concerned, seek medical advice.

Dispose in accordance with all federal, state and local regulations.

Application Rates and Frequencies in Canada: Follow vour professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.6 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 2.5 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N) Equivalent to 1.4 kg of product per 100 sq. meters.

| SUGGESTED SPREADER SETTINGS | | | |
|-----------------------------|-------------------------------|-------------------------------|--|
| Spreader | 0.43 kg N /100 m ² | 0.22 kg N /100 m ² | |
| Lebanon Turf | 41/2 | 31/2 | |
| Andersons Acc | uPro K | H | |
| Earthway Rotar | y 16 | 14 | |
| Gandy | 28 | 24 | |
| Lely | 6 II | 4 II | |
| Lesco (letter/ni | ımeric) i/17 | H/15 | |
| PennMulch HV |) L | J | |
| ProScape SS | L | J | |
| Spyker | 41/2 | 31/2 | |
| Vicon | 24 | 22 | |

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



22-35743

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com



For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight