LebanonTurf

Country Club Blended Fertilizer



FERTILIZER

25-0-15



GUARANTEED ANALYSIS		GUARANTEED MINIMUM ANALYSIS
Total Nitrogen (N) 3.9% Water Insoluble Nitrogen* 14.8% Urea Nitrogen 6.3% Other Water Soluble Nitrogen* Soluble Potash (K ₂ 0) Sulfur (S) 5.1% Combined Sulfur (S) Iron (Fe) 0.4% Water Soluble Iron (Fe)		Total Nitrogen (N)
Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate. *10.2% Slowly available nitrogen from Methylene Ureas. Chlorine (CI) not more than	2.0% U.S.A	Chlorine (Cl) (maximum)

25% Nitrogen' ıble Nitrogen

CANADA

NOTICE: This product contains the secondary micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Warning



Application Rates: To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.6 lbs per 1,000 sq. ft. or 157 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2 lbs

per 1,000 sq. it. or or ibs. per acre.			
SUGGES	SUGGESTED SPREADER SETTINGS		
Spreader 0.9#N	l/1,000 sq. ft.	0.5#N/1,000 sq. ft.	
Lebanon Turf	3	2	
Andersons AccuPro	F	D	
Earthway Rotary	12	10	
Gandy	27	23	
Lely	5½ II	4½ II	
Lesco (letter/numeric) I/17	G/15	
PennMulch HVO	J	E	
ProScape SS	J	E	
Spyker	3	2	
Vicon	22	16	
These settings were calibrated and field tested.			

However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Hazard Statements Causes mild skin irritation. Causes eye irritation on contact. May be harmful if inhaled repeatedly over long time periods. Suspected of causing cancer by prolonged/repeated inhalation.

Precautionary Statements If skin irritation occurs, wash with plenty of soap and water. Avoid breathing dust. Wash hands after handling.

Use appropriate personal protective equipment as required to avoid In case of eye contact, rinse cautiously with water for several minutes.

If irritation persists: seek medical attention. Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin, In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn. WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 1.7 kg of product per 100 sq. meters.

To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 0.9 kg of product per 100 sg. meters.

	3	
	SUGGESTED SPREADE	R SETTINGS
Spreader	0.43 kg #N/100 m ² .	0.22 kg #N/100 m ² .
Lebanon Turf	3	2
Andersons Ad	ccuPro F	D
Earthway Rot	tary 12	10
Gandy	27	23
Lely	5½ II	4½ II
Lesco (letter)	/numeric) I/17	G/15
PennMulch H	IVO J	E
ProScape SS	J	E
Spyker	3	2

Vicon 22 16
These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage



22-35760 | j/16

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042 800-233-0628 • (717) 273-1685



FERTILIZER



GUARANTEED ANALYSIS	
Total Nitrogen (N)	25%
3.9% Water Insoluble Nitrogen*	
14.8% Urea Nitrogen	
6.3% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ 0)	15%
Sulfur (S)	5.1%
Iron (Fe)	2.0%
0.4% Water Soluble Iron (Fe)	
Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate.	
*10.2% Slowly available nitrogen from Methylene Ureas.	
Chlorine (CI) not more than	2.0%

....

GUARANTEED WINIWOW ANALTSIS	
Total Nitrogen (N)	25%
3.9% Water Insoluble Nitrogen*	
14.8% Urea Nitrogen	
6.3% Other Water Soluble Nitrogen*	
Soluble Potash (K,0)	. 15%
Sulfur (S)	5.1%
5.1% Combined Sulfur (S)	
Iron (Fe) (Actual)	2.0%
0.4% Water Soluble Iron (Fe)	
Chlorine (CI) (maximum)	2.0%
Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sucrate.	
*10.2% Slowly available nitrogen from Methylene Ureas.	
	CANADA

CHARANTEED MINIMIM ANALYSIS

NOTICE: This product contains the secondary micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

U.S.A

Application Rates:

To feed at the rate of 0.9lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.6 lbs per 1,000 sq. ft. or 157 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2 lbs per 1,000 sq. ft. or 87 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
Spreader 0.9#N	/1,000 sq. ft.	0.5#N/1,000 sq. ft.
Lebanon Turf	3	2
Andersons AccuPro	F	D
Earthway Rotary	12	10
Gandy	27	23
Lely	51∕2 II	4½ II
Lesco (letter/numeric)	I/17	G/15
PennMulch HVO	J	E
ProScape SS	J	E
Spyker	3	2
Vicon	22	16
Those settings were colibrated and field tested		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Warning



Causes eye irritation on contact.

May be harmful if inhaled repeatedly over long time periods. Suspected of causing cancer by prolonged/repeated inhalation.

Precautionary Statements

Hazard Statements

If skin irritation occurs, wash with plenty of soap and water, Avoid breathing dust.

Wash hands after handling.

Use appropriate personal protective equipment as required to avoid breathing dust.

In case of eye contact, rinse cautiously with water for several minutes. If irritation persists: seek medical attention.

Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To apply at the rate of 0.43 kg N/100 sq. meters. Equivalent to 1.7 kg of product per 100 sq. meters.

To apply at the rate of 0.22 kg N/100 sq. meters. Equivalent to 0.9 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS		
Spreader 0.43 k	a #N/100 m².	0.22 kg #N/100 m².
Lebanon Turf	3	2
Andersons AccuPro	F	D
Earthway Rotary	12	10
Gandy	27	23
Lely	51/2 II	4½ II
Lesco (letter/numeric)	I/17	G/15
PennMulch HVO	J	E
ProScape SS	J	E
Spyker	3	2
Vicon	22	16

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



22-35760 i/16

FERTILIZER 29-0-10



MESA

GUARANTEED ANALYSIS		G
Total Nitrogen (N)	29%	Total Nitroger
3.5% Ammoniacal Nitrogen		3.5% Ami
2.2% Water Insoluble Nitrogen*		2.2% Wat
19.3% Urea Nitrogen		19.3% Ure
4.0% Other Water Soluble Nitrogen*		4.0% Oth
Soluble Potash (K ₂ 0)	12%	Soluble Potas
Sulfur (S)	4.2%	Sulfur (S)
5.7% Combined Sulfur (S)		5.7% Con
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of	Potash.	Chlorine (CI) ı
*6.2% Slowly Available Nitrogen from Methylene Ureas.		Derived From:
Chlorine (CI) not more than	9.0%	*6.2% Slowly
	U.S.A	

GUARANTEED MINIMUM ANALYSIS	
Total Nitrogen (N)	29%
3.5% Ammoniacal Nitrogen	
2.2% Water Insoluble Nitrogen*	
19.3% Urea Nitrogen	
4.0% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ 0)	12%
Sulfur (S)	4.2%
5.7% Combined Sulfur (S)	
Chlorine (CI) maximum	9.0%
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate o	f Potash.
*6.2% Slowly Available Nitrogen from Methylene Ureas.	
	CANADA

f savves anothe fields and landescuse Cuscific annihilation value and francescus

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.



Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 3.1 lbs per 1,000 sq. ft. or 135 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 1.7 lbs per 1,000 sq. ft. or 74 lbs. per acre.

SUGG	ESTED SPREADE	R SETTINGS
Spreader 0.	9# /1,000 sq. ft.	0.5# /1,000 sq. ft.
Lebanon Turf	41/2	31/2
Andersons AccuPr	o M	Н
Earthway Rotary	15	14
Gandy	30	24
Lely	6½ II	4 II
Lesco (letter/nume	eric) J/21	G/15
PennMulch HVO	Ĺ	i
ProScape SS	L	i
Spyker	41/2	31/2
Vicon	26	22
Those cottings w	ara palibrated a	nd field tected

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Warning

Hazard Statements

May be harmful if inhaled repeatedly over prolonged periods. Suspected of causing cancer by prolonged/repeated inhalation. Causes mild skin irritation.

May cause eye irritation on contact

Precautionary Statements

Avoid breathing dust.

Use appropriate personal protective equipment as required to avoid breathing dust.

If skin irritation occurs: Wash with soap and water.

Wash hands and exposed skin thoroughly after handling. If in eyes, rinse cautiously with water for several minutes. If eye irritation persists: seek medical attention.

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 1.5 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N) Equivalent to 0.9 kg of product per 100 sg. meters.

SUGGES	STED SPREAD	ER SETTINGS
Spreader 0.4	43# N/100 m²	0.22# N/100 m ²
Lebanon Turf	41/2	31/2
Andersons AccuPro	M	Н
Earthway Rotary	15	14
Gandy	30	24
Lely	6½ II	4 II
Lesco (letter/numeri	c) J/21	G/15
PennMulch HVO	Ĺ	i
ProScape SS	L	i
Spyker .	41/2	31/2
Vicon	26	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-29084

i/16

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com