

Lebanon Pro



STARTER FERTILIZER



18-24-12

GUARANTEED ANALYSIS		GUARANTEED MINIMUM ANALYSIS	
Total Nitrogen (N)	18%	Total Nitrogen (N)	18%
0.4% Water Insoluble Nitrogen*		0.4% Water Insoluble Nitrogen*	
17.6% Urea Nitrogen*		17.6% Urea Nitrogen*	
Available Phosphate (P ₂ O ₅)	24%	Available Phosphate (P ₂ O ₅)	24%
Soluble Potash (K ₂ O)	12%	Soluble Potash (K ₂ O)	12%
Derived From: Ammonium Phosphate, Biosolids, Polymer Coated Urea, Muriate of Potash.		Derived From: Ammonium Phosphate, Biosolids, Polymer Coated Urea, Muriate of Potash.	
*4.9% Slowly available Nitrogen from Polymer Coated Urea and biosolids.		Chlorine (Cl) maximum	
Chlorine (Cl) not more than	10.0%	*4.9% Slowly available Nitrogen from Polymer Coated Urea and biosolids.	
U.S.A		CANADA	

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:
To feed at the rate of 1 lb. Phosphate (P₂O₅) per 1,000 sq. ft., apply this product at 4.2 lbs per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.5 lb. Phosphate (P₂O₅) per 1,000 sq. ft., apply this product at 2.1 lbs per 1,000 sq. ft. or 91 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
Spreader	1 # P ₂ O ₅ /1,000 sq. ft.	0.5# P ₂ O ₅ /1,000 sq. ft.
Lebanon Turf	4½	4
Andersons AccuPro	1	H
Earthway Rotary	15	14
Gandy	25	24
Lely	5 II	5 II
Lesco (letter/numeric)	J/21	G/15
PennMulch HVO	K	J
ProScape SS	K	J
Spyker	4½	4
Vicon	23	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

- 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application;
- 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State;
- or 3) Re-establishing or repairing a turf area.

*In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

WARNING
Hazard Statements
Causes skin irritation.
Causes eye irritation.
May cause respiratory irritation.
May be harmful if inhaled repeatedly over prolonged periods.
Suspected of causing cancer by prolonged/repeated inhalation.



Precautionary Statements
Keep container tightly closed and store in a well-ventilated place.
Avoid breathing dust or spray mist.
Wash hands and exposed skin thoroughly after handling.
Use only outdoors or in a well-ventilated area.
Wear protective gloves/protective clothing/eye protection/face protection.
Avoid breathing dust.
Use appropriate personal protective equipment as required to avoid breathing dust.

IF ON SKIN: Gently wash with plenty of soap and water. Seek medical attention if irritation occurs and persists.
If in eyes: Rinse cautiously with water for several minutes. If eye irritation persists, seek medical attention.
Call a poison center or doctor physician if you feel unwell.
Take off contaminated clothing and wash before reuse.
Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE: To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:
To feed at the rate of 0.48 kg Phosphate (P₂O₅) Equivalent to 2.3 kg of product per 100 sq. meters.
To feed at the rate of 0.24 kg Phosphate (P₂O₅) Equivalent to 1.2 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS		
Spreader	0.48 kg P ₂ O ₅ /100 m ²	0.24 kg P ₂ O ₅ /100 m ²
Lebanon Turf	4½	4
Andersons AccuPro	1	H
Earthway Rotary	15	14
Gandy	25	24
Lely	5 II	5 II
Lesco (letter/numeric)	J/21	G/15
PennMulch HVO	K	J
ProScape SS	K	J
Spyker	4½	4
Vicon	23	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



0 88685 40408 5
F699 **22-40408** C/17

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

800-233-0628 • (717) 273-1685



Manufactured by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 1704
800-233-0628 • (717) 273-1685
Made in the U.S.A.
www.LebanonTurf.com

NET WT. 50 LBS. (22.7 KG.)

STARTER FERTILIZER



18-24-12

GUARANTEED ANALYSIS

Total Nitrogen (N) 18%
 0.4% Water Insoluble Nitrogen*
 17.6% Urea Nitrogen*
 Available Phosphate (P₂O₅) 24%
 Soluble Potash (K₂O) 12%

Derived From: Ammonium Phosphate, Biosolids, Polymer Coated Urea, Muriate of Potash.

*4.9% Slowly available Nitrogen from Polymer Coated Urea and biosolids.

Chlorine (Cl) not more than 10.0%

U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N) 18%
 0.4% Water Insoluble Nitrogen*
 17.6% Urea Nitrogen*
 Available Phosphate (P₂O₅) 24%
 Soluble Potash (K₂O) 12%

Derived From: Ammonium Phosphate, Biosolids, Polymer Coated Urea, Muriate of Potash.

Chlorine (Cl) maximum 10.0%

*4.9% Slowly available Nitrogen from Polymer Coated Urea and biosolids.

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 1 lb. Phosphate (P₂O₅) per 1,000 sq. ft., apply this product at 4.2 lbs per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.5 lb. Phosphate (P₂O₅) per 1,000 sq. ft., apply this product at 2.1 lbs per 1,000 sq. ft. or 91 lbs. per acre.

SUGGESTED SPREADER SETTINGS

Spreader	1 # P ₂ O ₅ /1,000 sq. ft.	0.5# P ₂ O ₅ /1,000 sq. ft.
Lebanon Turf	4½	4
Andersons AccuPro	i	H
Earthway Rotary	15	14
Gandy	25	24
Lely	5 II	5 II
Lesco (letter/numeric)	J/21	G/15
PennMulch HVO	K	J
ProScape SS	K	J
Spyker	4½	4
Vicon	23	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

- 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application;
 - 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State;
 - or 3) Re-establishing or repairing a turf area.
- *In Maryland, the laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

Hazard Statements

Causes skin irritation.
 Causes eye irritation.
 May cause respiratory irritation.
 May be harmful if inhaled repeatedly over prolonged periods.
 Suspected of causing cancer by prolonged/repeated inhalation.

WARNING



Precautionary Statements

Keep container tightly closed and store in a well-ventilated place.
 Avoid breathing dust or spray mist.
 Wash hands and exposed skin thoroughly after handling.
 Use only outdoors or in a well-ventilated area.
 Wear protective gloves/protective clothing/eye protection/face protection.
 Avoid breathing dust.
 Use appropriate personal protective equipment as required to avoid breathing dust.
 IF ON SKIN: Gently wash with plenty of soap and water. Seek medical attention if irritation occurs and persists.
 If in eyes: Rinse cautiously with water for several minutes. If eye irritation persists, seek medical attention.
 Call a poison center or doctor physician if you feel unwell.
 Take off contaminated clothing and wash before reuse.
 Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.48 kg Phosphate (P₂O₅) Equivalent to 2.3 kg of product per 100 sq. meters.

To feed at the rate of 0.24 kg Phosphate (P₂O₅) Equivalent to 1.2 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

Spreader	0.48 kg P ₂ O ₅ /100 m ²	0.24 kg P ₂ O ₅ /100 m ²
Lebanon Turf	4½	4
Andersons AccuPro	i	H
Earthway Rotary	15	14
Gandy	25	24
Lely	5 II	5 II
Lesco (letter/numeric)	J/21	G/15
PennMulch HVO	K	J
ProScape SS	K	J
Spyker	4½	4
Vicon	23	22

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-40408

C/17

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

800-233-0628 • (717) 273-1685



Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 1704

800-233-0628 • (717) 273-1685

Made in the U.S.A.

www.LebanonTurf.com

NET WT. 50 LBS. (22.7 KG.)