

LebanonTurf

Country Club[®] Blended Fertilizer

17-0-17



GUARANTEED ANALYSIS

Total Nitrogen (N)	17%
5.4% Water Insoluble Nitrogen*	
2.2% Urea Nitrogen	
9.4% Other Water Soluble Nitrogen *	
Soluble Potash (K₂O)	17%
Sulfur (S)	5.8%
5.8% Combined Sulfur (S)	
Iron (Fe).....	0.5%
0.005% Water Soluble Iron (Fe)	
Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sulfate.	
*14.8% Slowly Available Nitrogen from Methylene Ureas.	
Chlorine (Cl) not more than	2.0%

U.S.A

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	17%
5.4% Water Insoluble Nitrogen*	
2.2% Urea Nitrogen	
9.4% Other Water Soluble Nitrogen *	
Soluble Potash (K₂O)	17%
Sulfur (S)	5.8%
5.8% Combined Sulfur (S)	
Iron (Fe) (actual).....	0.5%
0.005% Water Soluble Iron (Fe)	
Chlorine (Cl) (maximum)	2.0%
Derived From: Methylene Ureas, Urea, Sulfate of Potash, Iron Sulfate.	
*14.8% Slowly Available Nitrogen from Methylene Ureas.	

CANADA

Notice: This product contains the micronutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after applications by sweeping or blowing. Do not wash off with water.

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 5.3 lbs per 1,000 sq. ft. or 230 lbs. per acre. To feed at the rate of ½ lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.9 lbs per 1,000 sq. ft. or 126 lbs. per acre.

SUGGESTED SPREADER SETTINGS

Spreader	0.9# N/1,000 sq. ft.	0.5 # N/1,000 sq. ft.
Lebanon Turf	4½	3
Andersons AccuPro	K	G
Earthway Rotary	15½	13
Gandy	26	23
Lely	5½ II	3½ II
Lesco (letter/numeric) i/17		F/12
PennMulch HVO	L	G
ProScape SS	L	G
Spyker	4½	3
Vicon	24	21

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

WARNING



Hazard Statements

Causes skin irritation.
Causes eye irritation.
May be harmful if inhaled.

Precautionary Statements

Wash hands and exposed skin thoroughly after handling.
Wear protective gloves/protective clothing/eye protection/face protection.
If in eyes, rinse with water and seek medical attention if symptoms persist.
Take off contaminated clothing and wash before reuse.
IF ON SKIN: Wash with plenty of soap and water.
P312: IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.
If skin irritation occurs: Get medical advice/ attention.
If eye irritation persists: Get medical advice/attention.
Keep out of reach of children
Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.
Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

Do not apply to frozen ground/turf.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 2.5 kg of product per 100 sq. meters.

To feed at the rate of 0.22 kg Nitrogen (N) Equivalent to 1.4 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

Spreader	0.43 kg N /100 m ²	0.22 kg N /100 m ²
Lebanon Turf	4½	3
Andersons AccuPro	K	G
Earthway Rotary	15½	13
Gandy	26	23
Lely	5½ II	3½ II
Lesco (letter/numeric) i/17		F/12
PennMulch HVO	L	G
ProScape SS	L	G
Spyker	4½	3
Vicon	24	21

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.regulatory-info-lebsea.com>

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042

800-233-0628 • (717) 273-1685

1000 Lbs. (453.6 kg) Net Weight

22-29117

F699

G/20