



# 0.125% StayGuard™

with FERTILIZER  
PREEMERGENCE WEED CONTROL


# 0-0-7

GUARANTEED ANALYSIS

Soluble Potash (K<sub>2</sub>O) ..... 7%

Derived from: Muriate of Potash.

Chlorine (Cl) not more than.....5%



**22-37262**

F699

### SUGGESTED SPREADER SETTINGS

Spreader	Pounds per Acre	Pounds per 1,000 sq. ft.	Pounds per Acre	Pounds per 1,000 sq. ft.
LebanonTurf	5	7½	Lesco (letter/Numeric) J/21	O/29
Andersons AccuPro N	N	T	PennMulch HVO	T
Earthway Rotary	17	25	ProScape SS	T
Gandy	29	37	Spyker	7½
Lely	6 II	8 II	Vicon (all models)	36

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

PROVIDES PREEMERGENCE WEED CONTROL IN

- LANDSCAPE GROWN ORNAMENTAL SHRUBS, TREES, CHRISTMAS TREES, GROUND COVERS, NON-BEARING FRUIT AND NUT TREES, VINES, AND BARE GROUND NON-CROP AREAS
- ESTABLISHED TURFGRASS

#### ACTIVE INGREDIENT:

Flumioxazin..... 0.125%

OTHER INGREDIENTS:..... 99.875%

TOTAL:..... 100.00%

\*2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione contains 2.5 lbs. flumioxazin per ton.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional precautionary statements and directions for use.

For Medical Emergencies Only, Call (877) 325-1840

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

EPA REG. NO. 228-749-961

StayGuard is a trademark of Nufarm Americas, Inc.

(L) (D)  
EPA Est. No. 961-PA-01:961-IL-01.

# Net Weight: 50 lbs (22.7 kg)

### FIRST AID

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear appropriate protective eye wear such as goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride, shoes, and socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply where runoff is likely to occur. **DO NOT** apply when weather conditions favor drift from treated areas. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to plants and must be used strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposures.

Under some conditions this product may have a potential to runoff to surface water or adjacent land. Where possible, use methods which reduce soil erosion, including no till, limited till and contour plowing; these methods also reduce pesticide runoff. Use of vegetation filter strips along rivers, creeks, streams, wetlands or on the downhill side of fields where runoff could occur will minimize water runoff is advised.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.

DO NOT enter or allow others to enter treated areas until dust has settled.

### RESISTANCE MANAGEMENT

For resistance management, this product is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

Rotate the use of this product or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective. Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed-management strategies for specific crops and weed biotypes.

For further information or to report suspected resistance, contact Lebanon Seaboard Corporation at (888) 208-1368.

### PRODUCT INFORMATION

This product is a selective preemergence herbicide that provides extended residual control of a wide variety of annual broadleaf and grassy weeds. This product controls weeds by inhibiting protoporphyrinogen oxidase, an essential enzyme required by plants for chlorophyll biosynthesis. Seedling weeds are controlled preemergence when exposed to sunlight following contact with the soil applied herbicide. This herbicide has limited postemergent activity against newly germinated seedlings of some weed species, but to ensure optimal control, apply before weed germination.

Most hardy woody ornamental shrubs, trees, ground covers and established turfgrass are not susceptible to this product, so long as the product is applied in accordance with the label. Use caution when applying to plants that are producing a new flush of growth because herbicide may injure tender, newly formed leaves if granules remain in contact with leaf surfaces. This product may cause leaf spotting on some species if granules remain in contact with leaf surfaces, especially on new flush or on foliage that is wet at the time of application. However, subsequent plant growth is generally unaffected. This product may injure some species if not applied in accordance with the label and may also injure ornamental species that are not listed on the label. Therefore, read all label precautions and restrictions before applying this product.

#### Restrictions and Limitations

- **DO NOT** apply more than 300 lb. (0.375 lb. ai) per acre in a single application.
- **DO NOT** apply more than 600 lb. (0.75 lb. ai) per acre per year. **DO NOT** apply more than 2 applications at 300 lb. (0.375 lb. ai) per acre or more than 3 applications at 200 lb. (0.25 lb. ai) per acre per year.
- **DO NOT** treat the same site more than one time every 8 weeks.
- **DO NOT** apply by air.
- **DO NOT** apply when winds are gusty or above 10 mph.
- **DO NOT** graze or feed livestock forage cut from treated areas.

#### Preemergence Application

This product provides effective preemergence control of weeds listed in Table 4. For optimal preemergence control, apply this product to clean, weed free soil or mulch before weed seeds germinate. Disturbing soil surfaces after application may reduce herbicide efficacy. Approximately 1/2 to 3/4 inch of rainfall, overhead sprinkler irrigation or hand irrigation is required to activate this product. Inadequate irrigation or rainfall following application may reduce effectiveness of this product. Weed control is most persistent under cooler temperatures.

#### Application Instructions

Apply this product with clean, well maintained granular application equipment. Calibrate application equipment prior to use according to manufacturer's guidelines. Check frequently to be sure equipment is working properly and distributing granules uniformly. Avoid skips and overlaps, as poor weed control or crop injury may occur.

- When applying to ornamentals, apply to dry foliage. To test for moisture, rub hands over plant foliage.
- When applying to turfgrass, ensure leaf surfaces are mostly dry.
- When possible, irrigate with 1/2 to 3/4 inch of water to activate the herbicide.
- If granules remain on foliage following application, lightly brushing or blowing plants will reduce the potential for plant injury.

#### Drift Management

**DO NOT** apply under circumstances where possible drift to unprotected persons or to food, forage or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption can occur.

- Make applications when the wind velocity favors on-target product deposition. Apply only when the wind speed is less than or equal to 10 mph. For all applications, wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.
- All application equipment must be properly maintained and calibrated using appropriate carriers.



Distributed by:

**Lebanon Seaboard Corporation**  
1600 E. Cumberland St. • Lebanon, PA 17042  
800-233-0628 • (717) 273-1685  
Made in the U.S.A.  
www.LebanonTurf.com



## CALIBRATION OF HERBICIDE APPLICATION EQUIPMENT

This product is a very active herbicide and can injure some plants if applied at excessive rates. The user is responsible for accurately calibrating herbicide application equipment so that this product is applied at the labeled rate of 200 to 300 lbs. (0.25-0.375 lbs. ai/A) per acre. The user assumes responsibility for any plant damage or other liability resulting from application of this product at excessive rates.

Table 1. Use Rate Table

Rate		
Pounds/Acre		Pounds/1,000 sq ft
200 (0.25 lb. ai/A)		4.6 (0.25 lb. ai/A)
300 (0.375 lb. ai/A)		6.9 (0.375 lb. ai/A)

### ORNAMENTAL PLANT INFORMATION

Most established woody ornamental plants are not susceptible to this product when the product is applied according to the label. Prior to applying on a commercial scale, test a small number of plants grown under similar growing conditions to ensure crop safety. Ensure that test plants are actively growing at the time of application and examined for 4 to 8 weeks for symptoms of plant injury. If test plants are dormant at the time of application, or enter dormancy within 8 weeks after application, continue the injury evaluation until after plants break dormancy in the spring.

Mild leaf spotting is the most common symptom of injury on ornamental plants, and usually appears within 2 to 4 days after application. Leaf spotting is most likely to occur when this product is applied to new flush that is wet at time of application. To minimize the risk of leaf spotting, completely read and follow the Application Instructions section of the label.

#### Treatment of Ornamental Plant Species Labeled as Sensitive

This product can cause significant injury to ornamental plants listed in Table 3. The user assumes responsibility for any plant damage that results from an application to plants listed as sensitive.

#### Treatment of Ornamental Plant Species NOT Listed on Table 4

Prior to applying this product on non-listed species, test a small number of plants grown under similar growing conditions to ensure crop safety. **The user assumes responsibility for any plant damage that results from an application to non-listed plant species.**

#### Treatment of Ornamental Plant Species with Wet Foliage

When this product is applied to dry plants, most granules will bounce or roll off the foliage before releasing herbicide residues. However, if this product is applied to plants that are wet from dew, irrigation water or rainfall, some granules may stick to foliage and release herbicide residues. Herbicide release onto wet foliage can injure some plant species.

#### Treatment of Recently Transplanted Liners

Under certain conditions, this product can cause significant injury to recently transplanted liners of some plant species that are not significant when more established (i.e. have formed a well-developed root system). Injury to recently transplanted liners is most likely when applied to smaller liners (less than one-gallon container size), is applied at above labeled rates, and is applied before potting media or soil has been settled by irrigation or rainfall. Therefore, **DO NOT apply to recently transplanted liners with root balls less than 4 inches in diameter until plants have been actively growing for at least 6 weeks after transplanting and have established a vigorous root system.**

#### Treatment of Herbaceous Plants, Tropical Shrubs and Foliage Plants

This product may severely injure many annual bedding plants (e.g. Petunia), and some herbaceous perennials (e.g. Hosta), tropical shrubs and foliage plants. Therefore, **DO NOT** apply this product over-the-top of herbaceous annual bedding plants, herbaceous perennials, tropical shrubs or foliage plants until after the user has confirmed the product’s safety on a small number of plants grown under standard growing conditions.

#### Treatment of Seedling Trees and Shrubs

This product may injure seedlings of some trees and shrubs if applied before seedlings have established a vigorous root system. Therefore, **DO NOT** apply to tree and shrub seedlings within one year after germination, unless the user has first confirmed the product’s safety on a small number of plants grown under standard growing conditions.

#### DIRECTIONS FOR USE IN LANDSCAPE GROWN ORNAMENTAL SHRUBS, TREES, CHRISTMAS TREES, GROUND COVERS, NON-BEARING FRUIT AND NUT TREES, VINES, AND BARE GROUND NON-CROP AREAS

In residential and commercial landscapes, this product must only be applied by commercial licensed applicators. This product applied at 200 to 300 lbs per acre (0.25 to 0.375 lb ai/A) (4.6 to 6.9 lbs per 1,000 sq ft) is an effective treatment for control of the weeds listed in Table 2 when applied to established landscape grown shrubs, trees, Christmas trees, ground covers, non-bearing fruit and nut trees and vines. Non-bearing fruit and nut trees including citrus, apples, pears, peaches, etc. are defined as plants that will not bear fruit for at least one year after application. This product may also be applied to maintain bare ground in non-crop areas in apartment complexes, gravel surfaces, ground mats, golf courses, office complexes, parks, parking areas, recreational sites, schools, sidewalks and other similar sites.

#### Precautions for Landscape Applications

This product can cause contact injury to foliage of landscape ornamentals if applied over-the-top of new flush, and especially if foliage is wet. To minimize crop injury potential, apply to dormant or non-actively growing landscape plants and avoid application to new flush.

#### Restrictions for Landscape Applications

- DO NOT** apply more than 300 lb (0.375 lb. ai) per acre in a single application.
- DO NOT** apply more than 600 lb (0.75 lb. ai) per acre per year.
- DO NOT** treat the same plants more than one time every 8 weeks.
- DO NOT** apply more than 2 applications at 300 lb. (0.375 lb. ai) per acre or more than 3 applications at 200 lb. (0.25 lb. ai) per acre per year.
- DO NOT** apply to foliage that is wet due to rainfall, irrigation or dew.
- DO NOT** apply over-the-top of ornamental plants not listed on this label until susceptibility is evaluated on a small number of plants grown under standard growing conditions.
- DO NOT** apply to plants that are under moisture stress or stress from insects, diseases, animals, winter injury, planting shock or any other stresses.
- DO NOT** apply to plants grown from seed until at least one year after seed germination, unless a product safety test is first conducted on a small number of plants.
- DO NOT** apply to plants whose leaves channel the herbicide granules to the leaf base.
- DO NOT** apply to herbaceous bedding plants or areas where bedding plants will be planted or transplanted within 3 months following application.
- DO NOT** apply to plants while in propagation.
- DO NOT** apply to liners that are in pots less than 4 inches in diameter.
- DO NOT** apply to recently transplanted liners with root balls that are less than 4” in diameter until liners have been actively growing for at least 6 weeks after transplanting and have established a vigorous root system.
- DO NOT** incorporate product into soil or potting media.
- DO NOT** apply in an enclosed greenhouse structure.
- DO NOT** graze or feed livestock forage cut from treated areas.

#### WEEDS CONTROLLED

When applied at 200 to 300 lbs. per acre (0.25 – 0.375 lbs. ai/A) before weeds germinate, this product will provide good to excellent preemergence control of the following grass and broadleaf weeds.

COMMON NAME	SCIENTIFIC NAME
Alyssum, Hoary	<i>Berteroa incana</i>
Amaranth	<i>Amaranthus palmeri</i>
Palmer	<i>Amaranthus spinosus</i>
Spiny	<i>Echinochloa crus-galli</i>
Barnyardgrass	<i>Desmodium tomentosum</i>
Beggarweed, Florida	<i>Cardamine hirsute</i>
Bittercress, Hairy	<i>Poa annua</i>
Bluegrass, Annual	<i>Medicago polymorpha</i>
Burclover, California	<i>Mollugo verticillate</i>
Carpetweed	
Chickweed	<i>Stellaria media</i>
Common	<i>Cerastium vulgatum</i>
Mouseear	
Crabgrass	<i>Digitaria sanguinalis</i>
Large	<i>Digitaria ischaemum</i>
Smooth	<i>Digitaria ciliaris</i>
Southern	<i>Croton glandulosus var septentrionalis</i>
Croton, Tropic	<i>Taraxacum officinale</i>
Dandelion	<i>Commelina benghalensis</i>
Dayflower, Benghal	<i>Eupatorium capillifolium</i>
Dogfennel	<i>Murdannia nudiflora</i>
Doveweed	<i>Eclipta prostrata</i>
Eclipta	<i>Erodium cicutarium</i>
Filaree, Redstem	
Footail	<i>Setaria verticillate</i>
Bristly	<i>Setaria faberi</i>
Giant	<i>Setaria viridis</i>
Green	<i>Setaria glauca</i>
Yellow	<i>Galinisoga ciliate</i>
Galinisoga, Hairy	<i>Geranium carolinianum</i>
Geranium, Carolina	<i>Eleusine indica</i>
Goosegrass, Indian	<i>Senecio vulgaris</i>
Groundsel, Common	<i>Lamium amplexicaule</i>
Henbit	<i>Zonza canadensis</i>
Horseweed	<i>Indigofera hirsute</i>
Indigo, Hairy	<i>Datura stramonium</i>
Jimsonweed	<i>Kochia scoparia</i>
Kochia	<i>Polygonum persicaria</i>
Lady’s Thumb	<i>Chenopodium album</i>
Lambsquarters, Common	<i>Marchantia polymorpha</i>
Liverwort, Common*	<i>Eragrostis diffusa</i>
Lovegrass, California	
Mallow	
Common	<i>Malva neglecta</i>
Little	<i>Malva parviflora</i>
Venice	<i>Hibiscus trionum</i>
Anthemis cotula	
Mayweed	
Morningglory	<i>Ipomoea hederacea var integruscula</i>
Entireleaf	<i>Ipomoea hederacea</i>
Ivyleaf	<i>Jacquemontia tamnifolia</i>
Smallflower	<i>Ipomoea purpurea</i>
Tall	<i>Bryum spp.</i>
Moss species	<i>Fatua villosa</i>
Mulberry Weed	<i>Panicum dichotomiflorum</i>
Nightshade	<i>Panicum texanum</i>
Black	<i>Alchemilla arvensis</i>
Eastern Black	<i>Sagina procumbens</i>
Panicum	<i>Thlaspi arvense</i>
Fall	<i>Phyllanthus tenellus</i>
Texas	
Parsley-Piert	<i>Amaranthus blitoides</i>
Pearlwort, Birdseye	<i>Amaranthus retroflexus</i>
Pennycress, Field	<i>Amaranthus hybridus</i>
Phyllanthus, Long-stalk	<i>Amaranthus albus</i>
Pigweed	<i>Matricaria matricarioides</i>
Prostrate	
Redroot	<i>Plantago major</i>
Smooth	<i>Plantago lanceolata</i>
Tumble	<i>Tribulus terrestris</i>
Pineapple weed	<i>Portulaca oleracea</i>
Plantain	<i>Richardia scabra</i>
Broadleaf	
Buckhorn	<i>Ambrosia artemisiifolia</i>
Puncturevine	<i>Ambrosia trifida</i>
Purslane, Common	<i>Calandrinia ciliata</i>
Pusley, Florida	<i>Barbarea vulgaris</i>
Ragweed	<i>Cassia occidentalis</i>
Common	<i>Sesbania exaltata</i>
Giant	<i>Capsella bursa-pastoris</i>
Redmaids	<i>Sida spinosa</i>
Rocket, Yellow	<i>Brachiaria platyphylla</i>
Senna, Coffee	<i>Polygonum pensylvanicum</i>
Sesbania, Hemp	<i>Sonchus oleraceus</i>
Shepherd’s-purse	
Sida, Prickly (Teaweed)	<i>Euphorbia prostrata</i>
Signalgrass	<i>Euphorbia maculata</i>
Smartweed, Pennsylvania	<i>Acanthospermum hispidum</i>
Southistle, Annual	<i>Crassocephalum crepidioides</i>
Surge	
Prostrate	
Spotted	
Starbur, Bristly	<i>Cirsium arvense</i>
Thickhead	<i>Salsola iberica</i>
Thistle	<i>Abutilon theophrasti</i>
Canada	
Russian	<i>Amaranthus rudis</i>
Velvetleaf	<i>Amaranthus tuberculatus</i>
Waterhemp	
Common	
Tall	<i>Solanum nigrum</i>
Woodsorrel, Common Yellow	<i>Solanum ptycanthum</i>

\*Will provide some postemergent control of liverwort which will be enhanced at higher temperatures and light levels.

#### SENSITIVE ORNAMENTAL SPECIES

Injury has been reported after application of this product to the plants listed in Table 3, especially when applied to small, recently transplanted liners. **DO NOT** apply this product to plants listed in Table 3. In some cases, only specific cultivars are listed because injury has not been reported on other cultivars of this species.

Table 3. SENSITIVE SPECIES

#### ORNAMENTAL SPECIES

COMMON NAME	SCIENTIFIC NAME
Butterfly Bush	<i>Buddleia davidii</i>
Clethra, Summersweet	<i>Clethra alnifolia</i>
Deutzia, Slender	<i>Deutzia gracilis “Nikko”</i>
Holly, Winterberry	<i>Ilex verticillata</i>
Hydrangea	<i>Hydrangea spp.</i>
Nandina	<i>Nandina domestica “Firepower”</i>
Plumbago, Cape	<i>Plumbago auriculata</i>
Spirea, Japanese	<i>Spiraea japonica</i>
Wax Myrtle, Southern (seedling)	<i>Myrica cerifera</i>

Injury other than temporary leaf spotting has not been observed on species listed in Tables 4-6 when this product was applied at the labeled rate to established plants with a well-developed root system. Some species listed in Tables 4-6 may be injured if applied before transplanted liners have established a well-developed root system (See Restrictions for Landscape Applications section in this label for complete details). This product has not been applied to all cultivars of listed species, or under all environmental conditions and cultural practices under which these species could be grown. Prior to applying on a commercial scale, test on a small number of plants grown under similar growing conditions to ensure crop safety.

Table 4. TREES INCLUDING CHRISTMAS TREES

COMMON NAME	SCIENTIFIC NAME
Arborvitae	<i>Thuja occidentalis</i>
American	<i>Thuja plicata</i>
Giant	<i>Thuja orientalis</i>
Oriental	
Ash	<i>Fraxinus pennsylvanica</i>
Green	<i>Fraxinus Americana</i>
White	<i>Betula spp.</i>
Birch	<i>Cedrus deodora</i>
Cedar, Deodora	<i>Malus spp.</i>
Crabapple, Flowering	<i>Populus deltoides</i>
Cottonwood	
Cypress	<i>Cupressus sempervirens</i>
Italian	<i>Chamaecyparis obtusa</i>
Tempehhof	<i>Cornus florida</i>
Dogwood, Florida	<i>Ulmus parvifolia</i>
Elm, Chinese	
Eucalyptus	<i>Eucalyptus cinerea</i>
Silver Dollar	<i>Eucalyptus camaldulensis</i>
Red Gum	<i>Ficus benjamina</i>
Ficus	
Fir	<i>Abies balsamea</i>
Balsam	<i>Pseudotsuga menzesii</i>
Douglas	<i>Abies fraseri</i>
Fraser	<i>Abies grandis</i>
Grand	<i>Abies koreana</i>
Korean	<i>Abies procera</i>
Noble	<i>Ginkgo biloba</i>
Ginkgo	
Hemlock	<i>Tsuga canadensis</i>
Eastern	<i>Tsuga mertensiana</i>
Mountain	<i>Tsuga heterophylla</i>
Western	<i>Syringa josikaea</i>
Lilac, Hungarian	<i>Magnolia lilliflora</i>
Magnolia, Lily	
Maple	<i>Acer ginnala</i>
Flame	<i>Abutilon hybridum</i>
Flowering	<i>Acer palmatum</i>
Japanese	<i>Acer rubrum</i>
Red	<i>Acer pensylvanicum</i>
Striped	<i>Morus alba</i>
Mulberry, White	
Oak	<i>Quercus ilicifolia</i>
Bear	<i>Quercus virginiana</i>
Live	<i>Quercus palustris</i>
Pin	<i>Quercus rubra</i>
Red	<i>Quercus phellos</i>
Willow	
Pine	<i>Pinus strobus</i>
Eastern White	<i>Pinus banksiana</i>
Jack	<i>Pinus bugeanae</i>
Lacebark	<i>Pinus taeda</i>
Loblolly	<i>Pinus contorta</i>
Longepole	<i>Pinus palustris</i>
Lodgepole	<i>Pinus mugo</i>
Mugo	<i>Pinus ponderosa</i>
Ponderosa	<i>Pinus clausa</i>
Sand	<i>Pinus sylvestris</i>
Scotch	<i>Pinus echinata</i>
Shortleaf	<i>Pinus elliotii</i>
Slash	<i>Pinus virginiana</i>
Virginia	<i>Prunus maritima</i>
Plum, Beach	<i>Podocarpus spp.</i>
Podocarpus	<i>Populus spp.</i>
Poplar	<i>Juniperus virginiana</i>
Red Cedar, Eastern	<i>Cercis canadensis</i>
Redbud	
Spruce	<i>Picea pungens</i>
Blue	<i>Picea glauca conica</i>
Dwarf Alberta	<i>Picea abies</i>
Norway	<i>Picea omorika</i>
Serbian	<i>Picea sitchensis</i>
Sitka	<i>Rhus typhina</i>
Sumac, Staghorn	<i>Liquidambar styraciflua</i>
Sweetgum, American	
Sycamore	<i>Platanus occidentalis</i>
American	<i>Platanus racemosa</i>
California	<i>Taxus cuspidata</i>
Yew, Japanese	

Table 5. SHRUBS AND GROUND COVERS	
COMMON NAME	SCIENTIFIC NAME
Abelia, Glossy	<i>Abelia x grandiflora</i>
Acacia	<i>Acacia spp.</i>
Anise, Purple	<i>Illicium floridanum</i>
Barberry	<i>Berberis spp.</i>
Japanese	<i>Berberis thunbergii</i>
William Penn	<i>Berberis x gladwynensis</i>
Bayberry	<i>Myrica pensylvanica</i>
Bottlebrush	<i>Callistemon spp.</i>
Boxwood, Littleleaf	<i>Buxus microphylla</i>
Camellia, Japanese	<i>Camellia japonica</i>
Cinquefoil, Shrubby	<i>Potentilla fruticosa</i>
Cotoneaster, Shrubby	<i>Cotoneaster dammeri</i>
Crepe Myrtle <sup>1</sup>	<i>Lagerstroemia indica spp.</i>
Elaeagnus	<i>Elaeagnus spp.</i>
English Ivy	<i>Hedera helix</i>
Euonymus, Winged <sup>2</sup>	<i>Euonymus alatus</i>
Firethorn, Scarlet	<i>Pyracantha coccinea</i>
Forsythia	
Border	<i>Forsythia x intermedia</i>
Weeping	<i>Forsythia suspensa</i>
White	<i>Abeliophyllum distichum</i>
Gardenia, Cape Jasmine	<i>Gardenia jasminoides</i>
Heath	<i>Erica cinerea</i>
Holly	
Chinese	<i>Ilex cornuta</i>
Inkberry	<i>Ilex glabra</i>
Japanese <sup>2</sup>	<i>Ilex crenata</i>
Meserve <sup>2</sup>	<i>Ilex x meserveae</i>
Nellie R. Stevens	<i>Ilex x Nellie R. Stevens</i>
Honeysuckle, Japanese	<i>Lonicera japonica</i>
Indian Hawthorn	<i>Rhaphiolepis indica</i>
Juniper	
Creeping	<i>Juniperus horizontalis</i>
Shore	<i>Juniperus conferta</i>
Singleseed	<i>Juniperus squamata</i>
Myoporum	<i>Myoporum parvifolium</i>
Oleander	<i>Nerium spp.</i>
Cedar Grape	<i>Mahonia spp.</i>
Pachysandra	<i>Pachysandra terminalis</i>
Photinia	<i>Photinia x fraseri</i>
Pieris, Japanese	<i>Pieris japonica</i>
Pittosporum, Japanese	<i>Pittosporum tobira</i>
Privet	
Chinese <sup>1</sup>	<i>Ligustrum sinense</i>
Japanese	<i>Ligustrum japonicum</i>
Pyracantha	<i>Pyracantha spp.</i>
Rhododendron, Catawba	<i>Rhododendron catawbiense</i>
Rose	<i>Rosa spp.</i>
Sweet Flag	<i>Acorus calamus</i>
Sweetspire, Virginia	<i>Itea virginica</i>
Tea Olive	<i>Osmanthus fragrans</i>
Viburnum	
Arrowwood	<i>Viburnum dentatum</i>
Pink Dawn	<i>Viburnum x bodnantense</i>
Sweet <sup>1</sup>	<i>Viburnum odoratissimum</i>

- Temporary injury reported at bud break on some cultivars.
- Injury reported following application to newly transplanted liners.

Table 6. NON-BEARING FRUIT AND NUT TREES AND VINES	
COMMON NAME	SCIENTIFIC NAME
Apple	<i>Malus spp.</i>
Blueberry, Huckleberry	<i>Vaccinium spp.</i>
Bramble	<i>Rubus spp.</i>
Cherry, Sweet	<i>Prunus avium</i>
Citrus Fruits	<i>Citrus spp.</i>
Grape	<i>Vitis spp.</i>
Olive	<i>Olea spp.</i>
Peach	<i>Prunus persica</i>
Pear	<i>Pyrus communis</i>
Prune	<i>Prunus spp.</i>
Stone Fruits	
Tree Nuts	
Walnut	<i>Juglans spp.</i>
Chestnut	<i>Castanea spp.</i>
Pecan	<i>Carya illionoinensis</i>
Pistachio	<i>Pistacia vera</i>
Almond	<i>Prunus dulcis</i>
Filbert	<i>Corylus maxima</i>

#### DIRECTIONS FOR USE ON ESTABLISHED TURFGRASS

This product may be applied as a single or split application to established Bermudagrass. This product provides preemergence control of annual bluegrass, goosegrass and other weeds found in Table 2. This product may be applied to turfgrass in such areas as apartment complexes, golf courses (naturalized areas, fairways, aprons, and roughs), cemeteries, sod farms, roadsides, sports fields, campgrounds, office complexes, parks, parking areas, recreational sites, schools, residential turf and other similar sites. Application of this product to turfgrass during spring green-up may cause some delay in green-up.

#### PREEMERGENCE APPLICATION

Apply 200 to 300 lbs. (0.25 – 0.375 lbs. ai/A) of this product per broadcast acre (4.6 to 6.9 lbs. per 1,000 sq. ft.) as a preemergence application prior to any weed germination. A higher application rate will provide longer and more consistent weed control. After application, if rain is not expected within 24 hours, irrigate the treated turf with up to 1/2” of water to obtain maximum weed control and quicker response from the fertilizer. When applying in areas with heavy weed pressure or in years with substantial rainfall following application of this product, a second application is recommended. However, DO NOT apply more than 600 lbs. (0.75 lb. ai/A) of this product per acre per year.

This product provides the best control of annual bluegrass when applied in the late summer/early fall prior to any plants germinating. Control may be less effective when applied in the winter after annual bluegrass has germinated. A second application (split application) of this product may be required to provide adequate season-long annual bluegrass control.

This product will provide best control of crabgrass, goosegrass and other summer annual weeds when applied in the late winter/early spring just prior to germination. Control will be less effective when applied in the spring after annual grass and broadleaf weeds have germinated. A second application (split application) of this product may be required to provide adequate season-long control of summer annual weeds.

#### USE AROUND BENTGRASS AND POA GREENS

This product has limited potential for lateral movement on level terrain but can potentially move down slope after excessive rainfall and affect sensitive turf species such as bentgrass, Poa annua and Poa trivialis. When applied upslope from bentgrass greens or Bermudagrass greens overseeded with Poa trivialis, allow an adequate buffer zone between greens and the treated area. If uncertain about the size of the buffer, 15 feet is suggested. Avoid application when heavy rain is imminent after application.

#### TOLERANT TURFGRASS SPECIES

If a desired turfgrass species is not listed in Tables 7[1][3], users are advised to evaluate the safety of this product in a small area under commercial growing conditions, and monitor plant response for four to six weeks for phytotoxicity. Testing this product in a small area will determine if this product can be used safely on a widespread basis.

TABLE 7 TOLERANT TURFGRASS SPECIES

COMMON NAME
Bermudagrass

#### RESTRICTIONS FOR APPLICATIONS TO TURFGRASS

- DO NOT** apply more than 300 lb. (0.375 lb. ai) per acre in a single application.
- DO NOT** apply more than 600 lb. (0.75 lb. ai) per acre per year.
- DO NOT** apply more than 2 applications at 300 lb. (0.375 lb. ai) per acre or more than 3 applications at 200 lb. (0.25 lb. ai) per acre per year
- DO NOT** treat the same treatment site more than one time every 8 weeks.
- DO NOT** apply to golf course putting greens or tees.
- DO NOT** apply to creeping bentgrass, or Dichondra.
- DO NOT** apply within 30 days prior to cutting or lifting sod.
- DO NOT** apply to seedling turfgrass until after receiving 2 mowings.

#### USE PRECAUTIONS FOR APPLICATIONS TO TURFGRASS