

Lebanon Pro



BRAND

FERTILIZER
WITH
.86% Pendimethalin
PREEMERGENT WEED CONTROL

13-0-5

Guaranteed Analysis

Total Nitrogen (N) 13%

13.0% Urea Nitrogen*

Soluble Potash (K₂O)..... 5%

Derived From: Urea, Muriate of Potash.

Chlorine (Cl) not more than..... 2.0%

* 3.25% Stabilized Urea Nitrogen with N-(n-butyl) thiophosphoric triamide.



F699 **22-40311**

g/14

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

SUGGESTED SPREADER SETTINGS

Spreader	4.0#/1,000 sq. ft.	5.3#/1,000 sq. ft.	Spreader	4.0#/1,000 sq. ft.	5.3#/1,000 sq. ft.
LebanonTurf	4%	5%	Lesco (letter/Numeric)	J/21	K/22
Andersons AccuPro	M	O	PennMulch HVO	N	O
Earthway Rotary	16	18	ProScape SS	N	O
Gandy	28	30	Spyker	4%	5%
Lely	6 ll	6½ ll	Vicon (all models)	26	30

Varied conditions will affect the application rates of spreaders; therefore the settings listed below are approximate. Age and conditions of spreader and walking speed can cause variation in application rate. Calibrate your spreader with each application. Close spreader hopper and adjust the setting using the suggested settings chart below. Fill spreader on a hard surface to eliminate the chance of spillage on the lawn. Close spreader hopper when stopping or turning.

FOR USE ON TURFGRASSES, ORNAMENTAL PLANTINGS, AND GROUND COVERS AS LISTED.

ACTIVE INGREDIENT:

Pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine 0.86%

OTHER INGREDIENTS: 99.14%

TOTAL: 100.00%

(50 pounds of this product contains 0.43 pound of the active ingredient, Pendimethalin)

EPA Reg. No. 19713-615-961

(L) (D)

EPA Est. No. 961-PA-01; 961-IL-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

See FIRST AID Below

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE, CALL CHEMTREC: 1-800-424-9300

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE): Some materials that are chemical-resistant to this product are any waterproof material (refer to Category A on an EPA chemical-resistance selection chart).

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the applicator at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep children and pets out of the treated area until dusts have settled. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in greenhouses, shadehouses or other enclosed structures. Not for use in commercial seed production.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and Restricted Entry Interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 24 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves made of any waterproof material and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the WPS for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter or allow others to enter the treated area until dusts have settled.

Failure to follow the directions for use and the precautions on this label may result in poor weed control, crop injury or illegal residues.

USE INFORMATION

LebanonPro Brand fertilizer with 0.86% Pendimethalin Preemergent Weed Control is a combination pre-emergent herbicide with fertilizer formulated to provide control of most annual grass weeds and certain broadleaf weeds found in Turfgrass grown in golf courses, lawns in residential, commercial establishments or grounds around military or other institutions, sod farms, parks, airports, school or cemeteries, prairiegrass areas, athletic fields or recreational areas, roadsides, pipe yards, industrial, unimproved Turf areas, and other sites where Turf may be maintained. This product is labeled for use on the following Turfgrass species: Kentucky bluegrass, Perennial ryegrass, Fine fescue, Tall fescue, Bentgrass (except putting greens and tees), *Poa annua* (except putting greens and tees), Bermudagrass, St. Augustinegrass, Bahiagrass, Centipedegrass, Zoysiagrass and blends of these grasses.

This product controls weeds as their seeds germinate, but will not control established weeds. Use a post-emergent herbicide, such as 2,4-D or MSMA, to control established weeds. Consult all herbicide label directions for proper use and precautions. Applied according to label directions and under normal growing conditions, this product will not cause Turf injury. Over application can result in Turf stand loss, Turf injury, or soil residues. Uneven application can decrease weed control or cause Turf injury.

Seeding diseases, cold weather, excessive moisture, high soil pH, soil salt concentration or drought can weaken Turfgrass and increase the possibility of Turf damage from this product.

Unusually cold, excessively wet or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

This product can be used around established Ornamentals and in established Wildflower plantings. This product may be used for hardwood and conifer regeneration on conservation reserve program land.

APPLICATION

Apply this product uniformly at the listed rate using a properly calibrated spreader (see table below for approximate settings). Avoid overlaps that will increase rates above those listed. The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the Turfgrass foliage and establish a pre-emergent herbicide barrier in the soil. If this product is not activated by rainfall or irrigation within 30 days of application, erratic weed control may result. Avoid contact of this product with porous surfaces such as driveways, stone or wood. Removing particles will help prevent staining.

This product provides pre-emergence control of the weed species listed. Applications can be made around and over the top of the Ornamentals and to the sites listed on this label.

MAXIMUM APPLICATION RATE BY USE SITES

Use Site	This Product/Ac. (Lbs.)	This Product/1,000 sq. ft. (Lbs.)	Pounds a.i./Ac.
Residential* and Sod farm turf	233	5.3	2.0
Golf course, Commercial and other Non-residential	349	8.0	3.0

* Residential is defined as Turf in any residential situation as well as home lawns, schools, parks and playgrounds.

TABLE OF EQUIVALENTS:

This Product Per Acre (Lbs.)	This Product Per 1,000 Sq. Ft. (Lbs.)
116	2.6
175	4.0
233	5.3
349	8.0

LebanonTurf

Distributed by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

U.S.A.

Net Weight 50 LBS. (22.7 kg)

FERTILIZER
WITH
.86% Pendimethalin
PREEMERGENT WEED CONTROL

13-0-5

Guaranteed Analysis

Total Nitrogen (N) 13%

13.0% Urea Nitrogen*

Soluble Potash (K₂O)..... 5%

Derived From: Urea, Muriate of Potash.

Chlorine (Cl) not more than..... 2.0%

* 3.25% Stabilized Urea Nitrogen with N-(n-butyl) thiophosphoric triamide.



F699 22-40311

g/14

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

SUGGESTED SPREADER SETTINGS

Spreader	4.0#/1,000 sq. ft.	5.3#/1,000 sq. ft.	Spreader	4.0#/1,000 sq. ft.	5.3#/1,000 sq. ft.
LebanonTurf	4 1/4	5 1/4	Lesco (letter/Numeric)	J/21	K/22
Andersons AccuPro	M	O	PennMulch HVO	N	O
Earthway Rotary	16	18	ProScape SS	N	O
Gandy	28	30	Spyker	4 3/4	5 1/4
Lely	6 ll	6 1/2 ll	Vicon (all models)	26	30

Varied conditions will affect the application rates of spreaders; therefore the settings listed below are approximate. Age and conditions of spreader and walking speed can cause variation in application rate. Calibrate your spreader with each application. Close spreader hopper and adjust the setting using the suggested settings chart below. Fill spreader on a hard surface to eliminate the chance of spillage on the lawn. Close spreader hopper when stopping or turning.

FOR USE ON TURFGRASSES, ORNAMENTAL PLANTINGS, AND GROUND COVERS AS LISTED.

ACTIVE INGREDIENT:

Pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine 0.86%

OTHER INGREDIENTS: 99.14%**TOTAL:** 100.00%

(50 pounds of this product contains 0.43 pound of the active ingredient, Pendimethalin)

EPA Reg. No. 19713-615-961

(L) (D)

EPA Est. No. 961-PA-01; 961-IL-01

KEEP OUT OF REACH OF CHILDREN**CAUTION**

See FIRST AID Below

FIRST AID**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE, CALL CHEMTREC: 1-800-424-9300

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

CAUTION: Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE): Some materials that are chemical-resistant to this product are any waterproof material (refer to Category A on an EPA chemical-resistance selection chart).

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in possession of the applicator at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep children and pets out of the treated area until dusts have settled. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in greenhouses, shadehouses or other enclosed structures. Not for use in commercial seed production.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and Restricted Entry Interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 24 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves made of any waterproof material and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the WPS for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter or allow others to enter the treated area until dusts have settled.

Failure to follow the directions for use and the precautions on this label may result in poor weed control, crop injury or illegal residues.

USE INFORMATION

LebanonPro Brand fertilizer with 0.86% Pendimethalin Preemergent Weed Control is a combination pre-emergent herbicide with fertilizer formulated to provide control of most annual grass weeds and certain broadleaf weeds found in Turfgrass grown in golf courses, lawns in residential, commercial establishments or grounds around military or other institutions, sod farms, parks, airports, school or cemeteries, prairiegrass areas, athletic fields or recreational areas, roadsides, pipe yards, industrial, unimproved Turf areas, and other sites where Turf may be maintained. This product is labeled for use on the following Turfgrass species: Kentucky bluegrass, Perennial ryegrass, Fine fescue, Tall fescue, Bentgrass (except putting greens and tees), *Poa annua* (except putting greens and tees), Bermudagrass, St. Augustinegrass, Bahiagrass, Centipedegrass, Zoysiagrass and blends of these grasses.

This product controls weeds as their seeds germinate, but will not control established weeds. Use a post-emergent herbicide, such as 2,4-D or MSMA, to control established weeds. Consult all herbicide label directions for proper use and precautions.

Applied according to label directions and under normal growing conditions, this product will not cause Turf injury. Over application can result in Turf stand loss, Turf injury, or soil residues. Uneven application can decrease weed control or cause Turf injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, soil salt concentration or drought can weaken Turfgrass and increase the possibility of Turf damage from this product.

Unusually cold, excessively wet or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

This product can be used around established Ornamentals and in established Wildflower plantings. This product may be used for hardwood and conifer regeneration on conservation reserve program land.

APPLICATION

Apply this product uniformly at the listed rate using a properly calibrated spreader (see table below for approximate settings). Avoid overlaps that will increase rates above those listed. The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the Turfgrass foliage and establish a pre-emergent herbicide barrier in the soil. If this product is not activated by rainfall or irrigation within 30 days of application, erratic weed control may result. Avoid contact of this product with porous surfaces such as driveways, stone or wood. Removing particles will help prevent staining.

This product provides pre-emergence control of the weed species listed. Applications can be made around and over the top of the Ornamentals and to the sites listed on this label.

MAXIMUM APPLICATION RATE BY USE SITES

Use Site	This Product/Ac. (Lbs.)	This Product/1,000 sq. ft. (Lbs.)	Pounds a.i./Ac.
Residential* and Sod farm turf	233	5.3	2.0
Golf course, Commercial and other Non-residential	349	8.0	3.0

* Residential is defined as Turf in any residential situation as well as home lawns, schools, parks and playgrounds.

TABLE OF EQUIVALENTS:

This Product Per Acre (Lbs.)	This Product Per 1,000 Sq. Ft. (Lbs.)
116	2.6
175	4.0
233	5.3
349	8.0



LebanonTurf

Distributed by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

U.S.A.

Net Weight 50 LBS. (22.7 kg)

GRASS WEEDS

Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-galli</i>	Junglerice	<i>Echinochloa colona</i>
Bluegrass, Annual	<i>Poa annua</i>	Lovegrass (from seed)	<i>Eragrostis spp.</i>
Crabgrass	<i>Digitaria spp.</i>	Panicum, Browntop	<i>Panicum fasciculatum</i>
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	Panicum, Fall	<i>Panicum dichotomiflorum</i>
Foxtail, Giant	<i>Setaria faberi</i>	Panicum, Texas	<i>Panicum texanum</i>
Foxtail, Green	<i>Setaria viridis</i>	Sandbur, Field	<i>Cenchrus incertus</i>
Foxtail, Yellow	<i>Setaria glauca</i>	Signalgrass	<i>Brachiaria platyphylla</i>
Goosegrass	<i>Eleusine indica</i>	Sprangletop, Mexican	<i>Leptochloa uninervis</i>
Itchgrass	<i>Rottboellia exaltata</i>	Sprangletop, Red	<i>Leptochloa filiformis</i>
Johnsongrass (from seed)	<i>Sorghum halepense</i>	Witchgrass	<i>Panicum capillare</i>
		Woolly Cupgrass	<i>Erochloa villosa</i>

BROADLEAF WEEDS

Common Name	Scientific Name	Common Name	Scientific Name
Burweed, Lawn	<i>Soliva pterospenna</i>	Pigweed	<i>Amaranthus spp.</i>
Carpetweed	<i>Mollugo verticillata</i>	Puncturevine	<i>Tribulus terrestris</i>
Chickweed, Common	<i>Stellaria media</i>	Purslane	<i>Portulaca oleracea</i>
Chickweed, Mouseear	<i>Cerastium vulgatum</i>	Pusley, Florida	<i>Richardia scabra</i>
Clover, Hop	<i>Trifolium procumbens</i>	Rocket, London	<i>Sisymbrium irio</i>
Cudweed	<i>Gnaphalium spp.</i>	Shepherdspurse	<i>Capsella bursa-pastoris</i>
Evening primrose	<i>Oenothera biennis</i>	Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Fiddleneck	<i>Amsinckia intermedia</i>	Speedwell, Corn	<i>Veronica arvensis</i>
Filaree	<i>Erodium spp.</i>	Spurge, Annual	<i>Euphorbia spp.</i>
Henbit	<i>Lamium amplexicaule</i>	Spurge, Prostrate	<i>Euphorbia numistrata</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>	Woodsorrel, Yellow	<i>Oxalis stricta</i>
Kochia	<i>Kochia scoparia</i>	Velvetleaf (Buttonweed)	<i>Abutilon theophrasti</i>
Lambsquarters	<i>Chenopodium album</i>		

TURFGRASSES

(Residential, Golf course, Commercial, Other Non-residential Turfgrass Uses) For Pre-emergence Weed Control

TURFGRASS SPECIES	WEEDS CONTROLLED	APPLICATION RATES*	COMMENTS
Cool Season Grasses Kentucky bluegrass, Perennial ryegrass, Fine fescue, Tall fescue	Crabgrass, Foxtail, <i>Poa annua</i> , Barnyardgrass, Fall panicum, <i>Oxalis</i> , Prostrate spurge, Purslane, Knotweed, Evening primrose, Hop clover	All Turf Uses: 175 to 233 pounds per acre as the initial application prior to weed germination in the Spring.	Apply a repeat application of 116 to 175 pounds per acre after 5 to 8 weeks for extended control or where heavy weed infestations are expected.
	Goosegrass	Residential** and Sod Turf Uses Only: 175 to 233 pounds per acre in the Spring prior to weed germination.	Apply a repeat application of 116 pounds per acre after 5 to 8 weeks if the lower rate was used initially or for extended Goosegrass control.
	Goosegrass	Golf course, Commercial, Non-residential Turf Uses Only: 175 to 349 pounds per acre in the Spring prior to weed germination.	Apply a repeat application of 116 pounds per acre after 5 to 8 weeks if the lower rate was used initially or for extended Goosegrass control.
	Cudweed, <i>Poa annua</i> , Chickweed, Lawn burweed, Henbit, Corn speedwell	All Turf Uses: 175 to 233 pounds per acre.	Apply in late Summer or early Fall prior to weed germination. Apply a repeat application of 175 to 233 pounds per acre after 5 to 8 weeks for extended <i>Poa annua</i> control.
Bentgrass or established <i>Poa annua</i> *** (one-half inch height or taller. Not for use on greens and tees.)	Crabgrass, Foxtail, <i>Poa annua</i> , Barnyardgrass, Fall panicum, <i>Oxalis</i> , Prostrate spurge, Purslane, Knotweed, Evening primrose, Hop clover	All Turf Uses (Non-Greens and Tees): 175 pounds per acre in the Spring prior to weed germination.	Apply a repeat application of 116 to 175 pounds per acre after 5 to 8 weeks where heavy weed infestations are expected.
Bentgrass or established <i>Poa annua</i> *** (one-half inch height or taller. Not for use on greens and tees.)	Goosegrass	All Turf Uses (Non-Greens and Tees): 175 pounds per acre.	Apply in the Spring prior to weed germination. Apply a repeat application of 175 pounds per acre after 5 to 8 weeks for extended control.
	Cudweed, <i>Poa annua</i> , Chickweed, Lawn burweed, Henbit, Corn speedwell	All Turf Uses (Non-Greens and Tees): 175 to 233 pounds per acre.	Apply in late Summer or early Fall prior to weed germination.
Warm Season Grasses Bermudagrass, St. Augustine grass, Bahagrass, Centipede grass, Zoysiagrass, Tall fescue, Buffalograss	Crabgrass, Foxtail, <i>Poa annua</i> , Barnyardgrass, Fall Panicum, <i>Oxalis</i> , Prostrate spurge, Purslane, Knotweed Evening primrose, Hop clover	Residential and Sod Farm Turf Uses Only: 175 to 233 pounds per acre prior to weed germination in the Spring.	Apply a repeat application of 116 to 175 pounds per acre after 5 to 8 weeks if necessary.
		Golf course, Commercial and Other Non-Residential Turf Uses Only: 175 to 349 pounds per acre in the Spring prior to weed germination.	
	Goosegrass	All Turf Uses (Non-Greens and Tees): 175 pounds per acre prior to weed germination in the Spring.	An additional application of 175 pounds per acre may be applied for extended Goosegrass control 8 weeks after the second application. Apply a repeat application of 175 to 233 pounds per acre after 5 to 8 weeks.
	Cudweed, Henbit, <i>Poa annua</i> , Chickweed, Lawn burweed, Corn speedwell	All Turf Uses: 175 to 233 pounds per acre.	Apply in late Summer or early Fall prior to weed germination. Apply a repeat application of 175 to 233 pounds per acre after 5 to 8 weeks for extended <i>Poa annua</i> control.

* DO NOT exceed a maximum rate of 233 pounds per acre per application for use on Residential Turfgrass and Sod farm Turfgrass.

DO NOT exceed a maximum rate of 349 pounds per acre per application for use on Golf course, Turfgrass, Commercial or other Non-residential Turfgrass.

** Residential is defined as Turf in any residential situation as well as home lawns, schools, parks and playgrounds.

*** Not for use on Bentgrass or *Poa annua* greens or tees.

USE PRECAUTIONS

- Avoid application when wind conditions may cause drift.
- Use on well-established Turfgrass with a dense and uniform stand. On newly planted areas, applications should not be made until the Turfgrass has filled in and has been mowed at least four times. On Turf that has been thinned or damaged due to Winter injury, excessive moisture, etc., allow for Turf recovery prior to making an application.
- Applications made to overseeded warm-season Turfgrasses may cause thinning or injury of the overseeded species.
- Do not use on Bentgrass or *Poa annua* greens and tees or injury may occur.
- Delay reseeding or Winter overseeding of treated Turfgrass for at least three (3) months following the last application of this product. Delay sprigging Turfgrass for five (5) months after application.

INDUSTRIAL (UNIMPROVED) TURF

Industrial or unimproved Turf areas often have a different spectrum of weeds to be controlled than those found in fine Turf as described elsewhere in this label. This product will control Annual grasses and Broadleaved weeds mentioned in that section of this label as well as the following weeds that might germinate in established grasses in rights-of-way, roadsides, construction sites, parks, substations, lots or similar areas: Crowfootgrass, Itchgrass, Johnsongrass (from seed), Junglerice, Lovegrass (from seed), Browntop Panicum, Texas Panicum, Field Sandbur, Signalgrass, Mexican Sprangletop, Red Sprangletop, Witchgrass, Woolly Cupgrass, Carpetweed, Fiddleneck, Filaree, Kochia, Lambsquarters, Pigweed, Puncturevine, Florida Pusley, London Rocket, Shepherdspurse, Pennsylvania Smartweed, Annual Spurge and Velvetleaf. Apply before weeds germinate. A post-emergence herbicide such as 2,4-D, MSMA or similar product is recommended to control established weeds.

LANDSCAPE AND GROUNDS MAINTENANCE

This product can be incorporated into landscape and ground maintenance programs to provide extended pre-emergence control of most annual grasses and certain broadleaf weeds. Areas to be treated include but are limited to established trees or Ornamentals in the landscape or in mulch beds; or other commercially maintained landscape areas such as borders, around statuary or monuments in parks or in parking areas. If emerged weeds are present at time of application, either shallow cultivate to destroy existing weeds or, where practical, remove by hand, or apply a post-emergence product labeled for such use prior to an application of this product. This product may be used before or after applications of herbicides registered for post-emergence use (i.e., Glyphosate) for the control of established weeds. Do not apply sprays containing Glyphosate over the top of desirable plants, and follow the label restrictions. A treatment of this product may be followed by any registered herbicide to control weeds not listed on this product's label.

ORNAMENTAL BULBS

This product may be applied for control of susceptible annual weeds in Ornamental bulbs listed on the label (Crocus, Daffodil (Narcissus), Gladiolus, Tulip, etc.). Apply this product prior to, during or after bulb emergence. If weeds have already germinated, use a labeled post-emergence herbicide to control emerged weeds.

WILDFLOWERS

This product may be applied for control of susceptible annual weeds in plantings of wildflowers listed on the label (Black-eyed Susan, California Poppy, Coreopsis, Oxeye Daisy, etc.). This product may be applied to established perennial wildflowers before emergence of weeds or wildflowers. For wildflowers being established from seed, apply this product after wildflowers have emerged but prior to weed germination. If weeds have already germinated, use a labeled post-emergence product to control emerged weeds. Due to the diversity of species and varieties which exist in areas where wildflowers are grown, the response to this product may vary greatly. Careful testing on desirable species is recommended to determine if area-wide applications can be made.

ORNAMENTAL PLANTINGS AND GROUND COVER

Use Precautions

Apply this product to established plantings. May be applied around and over the top of established plants listed below. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of this product with roots. Applications where soil and media surfaces are uniformly covered will result in best weed control and Ornamental tolerance. Plant only those desirable plant species listed on this label into soil treated the previous season with any Pendimethalin containing product or injury may occur.

TREES

Common Name	Scientific Name	Common Name	Scientific Name
Alder, European Black	<i>Alnus glutinosa</i>	Maple, Paper-bark	<i>Acer griseum</i>
Apple	<i>Malus spp.</i>	Maple, Norway	<i>Acer platanoides</i>
Arbortvitae, American	<i>Thuja occidentalis</i>	Maple, Japanese	<i>Acer palmatum</i>
Arbutus	<i>Arbutus spp.</i>	Maple, Red	<i>Acer rubrum</i>
Ash, Red	<i>Fraxinus pennsylvanica</i>	Maple, Sugar	<i>Acer saccharum</i>
Ash, White	<i>Fraxinus americana</i>	Nannyberry, Rusty	<i>Viburnum rufidulum</i>
Aspen, Bigtooth	<i>Populus grandidentata</i>	Oak, Chinquapin	<i>Quercus muehlenbergii</i>
Aspen, Quaking	<i>Populus tremuloides</i>	Oak, Live	<i>Quercus virginiana</i>
Basswood	<i>Tilia spp.</i>	Oak, Pin	<i>Quercus palustris</i>
Birch, European Weeping	<i>Betula pendula</i>	Oak, Red	<i>Quercus rubra</i>
Birch, River	<i>Betula nigra</i>	Oak, Swamp Chestnut	<i>Quercus michauxii</i>
Buckeye, Red	<i>Aesculus pavia</i>	Oak, Water	<i>Quercus nigra</i>
Cedar, White	<i>Thuja occidentalis</i>	Oak, White	<i>Quercus alba</i>
Chamaecyparis, Boulevard	<i>Chamaecyparis pisifera</i>	Oak, Willow	<i>Quercus phellos</i>
Cherry, Black	<i>Prunus serotina</i>	Olive	<i>Olea europaea</i>
Cherry, Choke	<i>Prunus virginiana</i>	Palm, Date	<i>Phoenix spp.</i>
Cherry, Kwanzan	<i>Prunus serrulata</i>	Palm, Fan	<i>Washingtonia spp.</i>
Cherry, Nanking	<i>Prunus tomentosa</i>	Palm, Pindo	<i>Butia spp.</i>
Cottonwood	<i>Populus deltoides</i>	Palm, Washington	<i>Washingtonia spp.</i>
Crabapple	<i>Malus spp.</i>	Peach	<i>Prunus persica</i>
Crepe Myrtle	<i>Lagerstroemia indica</i>	Pear, Bradford	<i>Pyrus calleryana 'Bradford'</i>
Cryptomeria, Japanese Cedar	<i>Cryptomeria japonica</i>	Pecan	<i>Carya illinoensis</i>
Cypress, Bald	<i>Taxodium distichum</i>	Pine, Austrian	<i>Pinus nigra</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	Pine, Italian Stone	<i>Pinus pinea</i>
Dogwood, Flowering	<i>Cornus florida</i>	Pine, Loblolly	<i>Pinus taeda</i>
Dogwood, Korean	<i>Cornus kousa</i>	Pine, Monterey	<i>Pinus radiata</i>
Dogwood, Silky	<i>Cornus amomum</i>	Pine, Red	<i>Pinus resinosa</i>
Dogwood, Shrub	<i>Cornus spp.</i>	Pine, Scotch	<i>Pinus sylvestris</i>
Elm	<i>Ulmus japonica</i>	Pine, Virginia	<i>Pinus virginiana</i>
Fir, Balsam	<i>Abies balsamiae</i>	Pine, White	<i>Pinus strobus</i>
Fir, Douglas	<i>Pseudotsuga menziesii</i>	Plum, Purple Leaf	<i>Prunus cerasifera</i>
Fir, Fraser	<i>Abies fraseri</i>	Poplar, Black	<i>Populus nigra</i>
Fir, White	<i>Abies concolor</i>	Redcedar, Eastern	<i>Juniperus virginiana</i>
Franklinia	<i>Franklinia spp.</i>	Redcedar, Western	<i>Thuja plicata</i>
Ginkgo	<i>Ginkgo biloba</i>	Red Ironbark	<i>Eucalyptus sideroxyylon 'Rosea'</i>
Gum, Black	<i>Nyssa sylvatica</i>	Redwood, Dawn	<i>Metasequoia glyptostroboides</i>
Gum, Sour	<i>Nyssa sylvatica</i>	Sequoia, Giant	<i>Sequoiadendron giganteum</i>
Haw, Black	<i>Viburnum prunifolium</i>	Serviceberry	<i>Amelanchier laevis</i>
Hawthorn	<i>Crataegus spp.</i>	Sourwood	<i>Oxydendrum arboreum</i>
Hemlock, Canada	<i>Tsuga canadensis</i>	Spruce, Colorado Blue	<i>Picea pungens</i>
Hemlock, Eastern	<i>Tsuga canadensis</i>	Spruce, Dwarf Alberta	<i>Picea glauca 'albertiana'</i>
Holly, American	<i>Ilex opaca</i>	Spruce, Norway	<i>Picea abies</i>
Honeylocust	<i>Gleditsia triacanthos</i>	Spruce, White	<i>Picea glauca</i>
Lilac, Common	<i>Syringa vulgaris</i>	Sweetgum	<i>Liquidambar styraciflua</i>
Lilac, Japanese Tree	<i>Syringa reticulata</i>	Sycamore	<i>Plantanus occidentalis</i>
Linden	<i>Tilia spp.</i>	Trachycarpus	<i>Trachycarpus spp.</i>
Magnolia, Saucer	<i>Magnolia soulangiana</i>	Tulip tree	<i>Liriodendron tulipifera</i>
Magnolia, Southern	<i>Magnolia grandiflora</i>	Walnut, Black	<i>Juglans nigra</i>
Magnolia, Star	<i>Magnolia stellata</i>	Willow, Weeping	<i>Salix babylonica</i>
Maidenhair Tree	<i>Ginkgo biloba</i>	Yellowwood	<i>Cladrastis lutea</i>

SHRUBS

Common Name	Scientific Name	Common Name	Scientific Name
Abelia, Glossy	<i>Abelia grandiflora</i>	Laurustinus	<i>Viburnum tinus</i>
Aucuba, Gold	<i>Aucuba japonica</i>	Lavender, English	<i>Lavandula angustifolia</i>
Azalea	<i>Rhododendron spp.</i>	Leucothoe	<i>Leucothoe fontanesiana</i>
Bamboo, Heavenly	<i>Nandina domestica</i>	Leucothoe, Coast	<i>Leucothoe axillaris</i>
Barberry	<i>Berberis gladwynesis</i>	Lilac, Cut-leaf	<i>Syringa laciniata</i>
Barberry, Japanese	<i>Berberis thunbergii</i>	Lily-of-the-Nile	<i>Agapanthus africanus</i>
Blue Indigo Bush	<i>Dalea greggii</i>	Mahonia	<i>Mahonia aquifolium</i>
Bottlebrush, Lemon	<i>Callistemon citrinus</i>	Mock Orange	<i>Pittosporum tobira</i>
Boxwood, Common	<i>Buxus sempervirens</i>	Myrtle, Compact	<i>Myrtus communis</i>
Boxwood, Japanese	<i>Buxus microphylla</i>	Myrtle, Wax	<i>Myrica cerifera</i>
Brittlebush	<i>Encelia farinosa</i>	Nandina	<i>Nandina domestica</i>
Buttonbush	<i>Cephalanthus occidentalis</i>	Oleander	<i>Nerium oleander</i>
Camellia	<i>Camellia japonica</i>	Oregon Grape	<i>Mahonia aquifolium</i>
Cape Jasmine	<i>Gardenia jasminoides</i>	Osmanthus	<i>Osmanthus fragrans</i>
Cassia, Feathery	<i>Cassia artemisioides</i>	Palm, European Fan	<i>Chamaerops humilis</i>
Cordyline	<i>Cordyline spp.</i>	Palm, Mediterranean Fan	<i>Chamaerops spp.</i>
Correa	<i>Correa spp.</i>	Phlox, Prickly	<i>Leptodactylon californicum</i>
Cotoneaster	<i>Cotoneaster apiculatus</i>	Photinia, Fraser	<i>Photinia x Fraseri</i>
Cotoneaster, Bearberry	<i>Cotoneaster dammeri</i>	Pieris, Japanese	<i>Pieris japonica</i>
Cotoneaster, Rock	<i>Cotoneaster horizontalis</i>	Pine, Mugo	<i>Pinus mugo</i>
Cypress, Italian	<i>Cupressus sempervirens</i>	Plum, Natal	<i>Carissa grandiflora</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	Privet, California	<i>Ligustrum ovalifolium</i>
Deutzia, Slender	<i>Deutzia gracilis</i>	Privet, Glossy	<i>Ligustrum lucidum</i>
Dogwood, Red Twig	<i>Cornus sericea</i>	Privet, Variegated	<i>Ligustrum sinensis</i>
Elaeagnus	<i>Elaeagnus ebbingei</i>	Privet, Waxleaf	<i>Ligustrum japonicum</i>
Escallonia	<i>Escallonia fradesii</i>	Pyracantha	<i>Pyracantha coccinea</i>
Euonymus	<i>Euonymus fortunei</i>	Quince, Flowering	<i>Chaenomeles japonica</i>
Euonymus, Golden	<i>Euonymus japonica</i>	Ranger, Texas	<i>Leucophyllum frutescens</i>
Euonymus, Winged	<i>Euonymus alata</i>	Redroot	<i>Ceanothus spp.</i>
Firethorn	<i>Pyracantha coccinea</i>	Rhododendron	<i>Rhododendron spp.</i>
Forsythia, Border	<i>Forsythia intermedia</i>	Robira	<i>Pittosporum tobira</i>
Fragrant Olive	<i>Osmanthus fragrans</i>	Rose	<i>Rosa spp.</i>
Fuschia, California	<i>Zauschneria californica</i>	Spice Plant	<i>Illicium parviflorum</i>
Gardenia	<i>Gardenia jasminoides</i>	Spiraea	<i>Spiraea vanhouttei</i>
Hawthorne, Indian	<i>Raphiolepis indica</i>	Spiraea, Anthony Waterer	<i>Spiraea x bumalda</i>
Hibiscus	<i>Hibiscus syriacus</i>	Spiraea, Japanese	<i>Spiraea japonica</i>
Holly, Chinese	<i>Ilex cornuta</i>	Sweet Bay	<i>Laurus nobilis</i>
Holly, Japanese	<i>Ilex crenata</i>	Trumpet Bush	<i>Tecoma stans</i>
Holly, Fosters	<i>Ilex attenuata 'Fosteri'</i>	Verbena, Lemon	<i>Aloysia triphylla</i>
Holly, Savannah	<i>Ilex attenuata</i>	Viburnum	<i>Viburnum suspensum</i>
Holly, Yaupon	<i>Ilex vomitoria</i>	Vitex	<i>Vites spp.</i>
Honeysuckle Bush	<i>Diervilla lonicera</i>	Weigela	<i>Weigela florida</i>
Hopseed Bush	<i>Dodonaea viscosa</i>	Wild Lilac	<i>Ceanothus spp.</i>
Hopbush	<i>Dodonaea viscosa</i>	Wisteria	<i>Wisteria spp.</i>
Hydrangea	<i>Hydrangea macrophylla</i>	Xylosma	<i>Xylosma congestum</i>
Juniper	<i>Juniperus spp.</i>	Yellowbells	<i>Tecoma stans</i>
Juniper, Chinese	<i>Juniperus chinensis v. pfitzer</i>	Yew	<i>Taxus media</i>
Juniper, Shore	<i>Juniperus conferta</i>	Yew, Japanese	<i>Taxus cuspidata</i>
Juniper, Trailing	<i>Juniperus horizontalis</i>	Yew, Southern	<i>Podocarpus macrophyllum</i>
Laurel, Cherry	<i>Prunus laurocerasus</i>	Yucca, Adam's Needle	<i>Yucca filamentosa</i>
Laurel, Mountain	<i>Kalmia latifolia</i>	Yucca, Weeping	<i>Yucca pendula</i>
Laurel, Otto Luyken	<i>Prunus laurocerasus</i>		
Laurel, Schipka	<i>Prunus schipkanesis</i>		

GROUND COVERS

Common Name	Scientific Name	Common Name	Scientific Name
Ajuga	<i>Ajuga reptans</i>	Morningglory	<i>Convolvulus spp.</i>
Baby Sun Rose	<i>Aptenia cordifolia</i>	Myoporum	<i>Myoporum parvifolium</i>
Beach Strawberry	<i>Fragaria chiloensis</i>	Pachysandra	<i>Pachysandra terminalis</i>
Capeweed	<i>Arctotheca calendula</i>	Potentilla	<i>Potentilla fruticosa</i>
Cinquefoil, Spring	<i>Potentilla verna</i>	Periwinkle, Common	<i>Vinca minor</i>
Coyotebrush, Dwarf	<i>Baccharis pitarularis</i>	Red Apple	<i>Aptenia cordifolia</i>
Daisy, Trailing African	<i>Osteospermum fruticosum</i>	Rosemary	<i>Rosemarinus officinalis</i>
Dymondia	<i>Dymondia margaretae</i>	Rose-Of-Sharon	<i>Hypericum calycinum</i>
Gazania	<i>Gazania splendens</i>	Sand Strawberry	<i>Fragaria chiloensis</i>
Iceplant, Large Leaf	<i>Carpobrotus edulis</i>	Sedum	<i>Sedum spurium</i>
Ivy, English	<i>Hedera helix</i>	St. Johnswort Creeping	<i>Hypericum calycinum</i>
Ivy, Geranium	<i>Pelargonium peltatum</i>	Stoncrop	<i>Sedum spurium</i>
Jasmine, Asiatic	<i>Trachelospermum asiaticum</i>	Verbena, Peruvian	<i>Verbena peruviana</i>
Jasmine, Primrose	<i>Jasminum mesnyi</i>	Vervain	<i>Verbena peruviana</i>
Jessamine, Carolina	<i>Gelsemium sempervirens</i>	Vetch, Crown	<i>Vicia sativa</i>
Manzanita, Bearberry	<i>Arctostaphylos uva-ursi</i>	Vinca or Greater Periwinkle	<i>Vinca major</i>
Miscanthus	<i>Miscanthus spp.</i>	Wintercreeper	<i>Euonymus fortunei</i>
Mondograss	<i>Ophiopogon japonica</i>		

PERENNIALS

Common Name	Scientific Name	Common Name	Scientific Name
Acacia	Acacia redolens	Gayfeather	Liatris spp.
Asparagus	Asparagus spp.	Gaillardia	Gaillardia pulchella
Aster, New York	Aster novi-belgii	Geum	Geum spp.
Aster, Stokes	Stokesia laevis	Gladiolus	Gladiolus spp.
Astilbe (False Spirea)	Astilbe spp.	Heather, Dwarf	Calluna vulgaris
Babys Breath	Gypsophila elegans	Hosta	Hosta spp.
Beard-Tongue	Penstemon spp.	Hyssop, Sunset	Agastache rupestris
Bellflower	Campanula spp.	Indian Blanket	Gaillardia pulchella
Bird of Paradise	Caesalpinia pulcherrima	Lantana, Weeping	Lantana montevidensis
Black-eyed Susan	Rudbeckia hirta	Leopards Bane	Doronicum cordatum
Blanket Flower	Gaillardia x grandiflora	Lily	Lilium spp.
Bleeding Heart	Dicentra spectabilis	Liriope, Big Blue	Liriope muscari
Butterfly Weed	Asclepias tuberosa	Liriope, Creeping	Liriope spicata
California Poppy	Eschscholzia californica	Lirope, Variegated	Liriope muscari
Calla Lily	Zantedeschia aethiopica	Loosestrife, Garden	Lysimachia punctata
Canna, Common Garden	Canna generalis 'Lucifer'	Moonbeam	Coreopsis verticillata
Carex	Carex spp.	Montbretia	Crococsmia crocosmiiflora
Cast Iron Plant	Aspidistra elatior	Nightshade	Solanum spp.
Chincherinchee	Ornithogalum thyrsoides	Orchid, Peacock	Acidanthera bicolor
Clover, Crimson	Trifolium incarnatum	Oxeye Daisy	Chrysanthemum leucanthemum
Columbine	Aquilegia 'McKana Giant'	Peony, Chinese	Paeonia lactiflora
Coreopsis	Coreopsis lanceolata	Purple Coneflower	Echinacea purpurea
Crinum Lily	Crinum spp.	Purple Loosestrife	Lythrum virgatum
Crocus	Crocus spp.	Rosemary	Rosmarinus officinalis
Daffodil	Narcissus spp.	Ruscus, Israeli	Ruscus hypophyllum
Daylily	Hemerocallis spp.	Sage, Russian	Perovskia atriplicifolia
False dragonhead	Physostegia virginiana	Sedge	Carex spp.
Fairy Duster	Calliandra eriophylla	Shasta Daisy	Chrysanthemum x superbum
Fern, Asparagus	Asparagus officinalis	Sweet Flag	Acorus calamus
Fern, Tree	Asparagus virgatus	Tickseed	Coreopsis lanceolata
Fern, Boston	Nephrolepis exaltata	Texas Bluebonnet	Lupinus texensis
Fern, Leatherleaf*	Rumohra adiantiformis	Tulip	Tulipa spp.
Fortnight Lily	Moraea spp.	Twinspur	Diascascia barbaerae
Foxglove, Common	Digitalis purpurea	Wonder Flower	Ornithogalum thyrsoides
Freesia	Freesia x hybrida	Yarrow	Achillea millefolium
Fuschia, Arizona	Zauschneria californica	Zephyr Lily	Zephyranthes spp.

* Applications of this product to immature ferns (during periods of new growth of fronds) may result in some injury.

ORNAMENTAL GRASSES

Common Name	Scientific Name	Common Name	Scientific Name
Beach Grass	<i>Ammophila breviligulata</i>	Reed Canary Grass	<i>Phalaris arundinacea</i>
Fescue, Blue	<i>Festuca ovina</i>	Reed, Giant	<i>Arundo spp.</i>
Fescue, Sheep	<i>Festuca ovina</i>	Ribbon Grass	<i>Phalaris arundinacea</i>
Fountain Grass	<i>Pennisetum setaceum</i>	Tufted Hair Grass	<i>Deschampsia caespitosa</i>
Pampas Grass	<i>Cortaderia selloana</i>		

BEDDING PLANTS*

Common Name	Scientific Name	Common Name	Scientific Name
Ageratum	<i>Ageratum houstonianum</i>	Impatiens	<i>Impatiens spp.</i>
Alyssum	<i>Alyssum saxatile</i>	Kale, Ornamental	<i>Brassica napus</i>
Anemone, Poppy-flowered	<i>Anemone coronaria</i>	Marigold, African	<i>Tagetes erecta</i>
Artemesia	<i>Artemesia spp.</i>	Moss Rose	<i>Portulaca grandiflora</i>
Balloonflower	<i>Platycodon grandiflorum</i>	Mum, Garden	<i>Chrysanthemum spp.</i>
Begonia	<i>Begonia spp.</i>	Pansy, Garden	<i>Viola x Wittrockiana</i>
Cabbage, Ornamental	<i>Brassica oleracea</i>	Periwinkle, Rose	<i>Catharanthus roseus or Vinca rosea</i>
Caladium	<i>Caladium spp.</i>	Petunia	<i>Petunia spp.</i>
China Aster	<i>Callistephus chinensis</i>	Plumosa Cockscornb	<i>Celosia cristata</i>
Crococsmia, Montebretia	<i>Crococsmia x crocosmiiflora</i>	Portulaca	<i>Portulaca grandiflora</i>
Dahlia	<i>Dahlia spp.</i>	Salvia	<i>Salvia splendens</i>
Dianthus	<i>Dianthus barbatus</i>	Snapdragon	<i>Antirrhinum majus</i>
Dusty Miller	<i>Senecio cineraria</i>	Statice	<i>Limonium spp.</i>
Gazania, Treasure Flower	<i>Gazania rigens</i>	Sweet William	<i>Dianthus barbatus</i>
Gazania, Trailing	<i>Gazania rigens leucolaena</i>	Vinca or Periwinkle	<i>Catharanthus roseus</i>
Geranium	<i>Geranium spp.</i>	Zinnia	<i>Zinnia spp.</i>
Gloxinia	<i>Gloxinia simningia</i>		

* After planting bedding plants and before applying this product, area to be treated should be watered sufficiently to pack soil around roots.

DO NOT treat plants grown for food or feed. DO NOT use treated plants for food or feed.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited. Do not reuse empty containers.

PESTICIDE STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing. Do not store above 120°F.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable Container (flexible-bag-all weights): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

For Residential/Household Use:

PESTICIDE DISPOSAL: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If empty,** place in trash or offer for recycling if available. **If partially filled,** call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, the User assumes responsibility for any crop damage or other liability when this product is used on plant species not listed on this label.

To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.