EBANON

FERTILIZER

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
6.0% Water Insoluble Nitrogen* 2.2% Urea Nitrogen	
9.6% Other Water Soluble Nitrogen*	1.00/
Soluble Potash (K ₂ O) Sulfur (S)	10% 6.0%
Derived From: Ammonium Sulfate, Methylene Ureas, Sulfate of Potash.	
*15.6% Slowly available Nitrogen from Methylene	Jreas.
Oblavina (OI) nat mana than	0.00/

Chlorine (CI) not more than 2.0%

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	20%
Soluble Potash (K ₂ O)	10%
Sulfur (S)6.0% Combined Sulfur (S)	6.0%
Chlorine (CI) (maximum)	2.0%
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Sulfate of Potash. *15.6% Slowly available Nitrogen from Methylene Ureas.	

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates: To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.5 lbs per 1,000 sq. ft. or 196 lbs. per acre. To feed at the rate of 0.5 lb.

Nitrogen (N) per 1,000 sq. ft., apply this product at 2.5

lbs per 1,000 sq. ft. or 109 lbs. per acre.

ina per 1,000 aq. it.	ins per 1,000 sq. it. or 103 ins. per acre.		
SUGGEST	SUGGESTED SPREADER SETTINGS		
Spreader 0.9# N	/1,000 sq. ft	t. 0.5# N /1,000 sq. ft	
Lebanon Turf	5	4	
Andersons AccuPro	N	i	
Earthway Rotary	17	14	
Gandy	29	25	
Lely	6 II	4½ II	
Lesco (letter/numeric)	J/21	H/16	
PennMulch HVO	M	J	
ProScape SS	M	J	
Spyker	5	4	
Vicon	28	23	
These settings were	calibrated	and field tested.	

However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage. Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of

contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Use in accordance with recommendations of a qualified individual or

institution, such as, but not limited to a certified crop advisor, agronomist university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to

WARRANTY

NOTE To the extent consistent with applicable law. Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

WARNING **Hazard Statements**

Causes skin irritation. Causes eye irritation. May be harmful if inhaled. Suspected of causing cancer.

Precautionary Statements for handling:

Do not handle until all safety precautions have been read and

Avoid breathing dust/fume/gas/mist/vapours/spray. Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.

Use personal protective equipment as required. If in eyes, rinse with water and seek medical attention if symptoms

Take off contaminated clothing and wash before reuse. IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Call » POISON CENTER or doctor/physician if you feel

If skin irritation occurs: Get medical advice/ attention. If eye irritation persists: Get medical advice/attention.

federal, state and local regulations.

Dispose of contents/container properly as required by law for waste pesticides.

If exposed or concerned, seek medical advice. Keep out of reach of children Precautionary Statements for disposal - Dispose in accordance with all

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Do not apply to frozen ground.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m2, applied every 4 to 8 weeks during the local growing season.

Application Rates: To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 3.5 kg of product per 100 sq. meters.

SUGGESTED SPREADER SETTINGS

To feed at the rate of 0.24 kg Nitrogen (N) Equivalent to 1.0 kg of product per 100 sq. meters.

Spreader	0.43 kg N/100 m ²	0.24 kg# N/100
Lebanon Turf	5	4
Andersons Accu	Pro N	i
Earthway Rotary	17	14
Gandy	29	25
Lelv	6 II	41/2 II
Lesco (letter/nu	meric) J/21	H/16
PennMulch HVO		J
ProScape SS	M	J
Spyker	5	4
Vicon	28	23

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage





Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

> Made in the U.S.A. www.LebanonTurf.com

NET WEIGHT LBS.(22.7 Kg)

FERTILIZER 20-0-10 AESA EXPORASIUM SULFATE

GUARANTEED ANALYSIS

Total Nitrogen (N)	
6.0% Combined Sulfur (S) Derived From: Ammonium Sulfate, Methylene Ureas,	
Sulfate of Potash. *15.6% Slowly available Nitrogen from Methylene	Ureas.
Chlorine (CI) not more than	
0	U.S.A

GUARANTEED MINIMUM ANALYSIS

2.2% Ammoniacal Nitrogen 6.0% Water Insoluble Nitrogen*	20%
2.2% Urea Nitrogen 9.6% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	10% 6.0%
Chlorine (CI) (maximum)	2.0%
Derived From: Ammonium Sulfate, Methylene Ureas, Sulfate of Potash.	Urea,

*15.6% Slowly available Nitrogen from Methylene Ureas.

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 0.9 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 4.5 lbs per 1,000 sq. ft. or 196 lbs. per acre. To feed at the rate of 0.5 lb. Nitrogen (N) per 1,000 sq. ft., apply this product at 2.5 lbs per 1,000 sq. ft. or 109 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
/1,000	sq. ft. 0.5# N /1,000 sq. ft.	
5	4	
N	i	
17	14	
29	25	
6 II	4 ½ Ⅱ	
J/21	H/16	
M	J	
M	J	
5	4	
28	23	
	/1,000 5 N 17 29 6 II J/21 M M	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

WARNING



Precautionary Statements for handling:

Do not handle until all safety precautions have been read and understood.

Avoid breathing dust/fume/gas/mist/vapours/spray.

Wash hands and exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.

Use personal protective equipment as required.

If in eyes, rinse with water and seek medical attention if symptoms persist.

Take off contaminated clothing and wash before reuse.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell

If skin irritation occurs: Get medical advice/ attention.

If eye irritation persists: Get medical advice/attention.

Store locked up.

Dispose of contents/container properly as required by law for waste pesticides.

If exposed or concerned, seek medical advice.

Keep out of reach of children

Precautionary Statements for disposal - Dispose in accordance with all federal, state and local regulations.

Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

Do not apply to frozen ground.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.8 Kg for 100 m2, applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.43 kg Nitrogen (N) Equivalent to 3.5 kg of product per 100 sq. meters.

To feed at the rate of 0.24 kg Nitrogen (N) Equivalent to 1.0 kg of product per 100 sq. meters.

Equivalent to 110 kg of product por 100 cq. motorer		
SUGGESTED SPREADER SETTINGS		
Spreader	0.43 kg N/100 m ²	0.24 kg# N/100 m ²
Lebanon Turf	5	4
Andersons Accul	Pro N	i
Earthway Rotary	17	14
Gandy	29	25
Lely	6 II	4½ II
Lesco (letter/nur	neric) J/21	H/16
PennMulch HVO	M	J
ProScape SS	M	J
Spyker	5	4
Vicon	28	23

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



F699

22-53717



Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

800-233-0628 • (717) 273-1685

Made in the U.S.A.

www.LebanonTurf.com

NET WEIGHT 50 LBS. (22.7 Kg)