

FERTILIZER WITH TEAM®PRO 0.86%

PREEMERGENCE WEED CONTROL

16-0-8 WITH MESA.

GUARANTEED ANALYSIS	
Total Nitrogen (N)	16%
2.4% Ammoniacal Nitrogen	
1.4% Water Insoluble Nitrogen*	
9.7% Urea Nitrogen	
2.5% Other Water Soluble Nitrogen*	
Soluble Potash (K2O)	8%
Sulfur (S)	2.8%
2.8% Combined Sulfur (S)	
Iron (Fe)	2.0%
0.2% Water Soluble Iron (Fe)	
Derived from: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of	
Potash, Ferric Oxide, Ferrous Sulfate.	



F699	22-53853
	H/21

Spreader Settings as a Guide for Calibration							
Brand and Model of Spreader	174#/acre	233#/acre	Spreader	174#/acre	233#/acre		
LebanonTurf Andersons AccuPro Earthway Rotary Gandy Lely	4¾ M 16 28 6 II	5¼ 0 19 31 6½ II	Lesco (letter/Numeric) PennMulch HVO ProScape SS Spyker Vicon (all models)	J/21 M M 43/4 26	K/22 0 0 5¼ 30		

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, Poa annua and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals.

ACTIVE INGREDIENTS:

Chlorine (CI) not more than......

*3.9% Slowly Available Nitrogen from Methylene Ureas.

benefin: <i>N</i> -butyl-N-ethyl- α , α , α -trifluoro-2,6-dinitro- p -toluidine	0.43%
trifluralin: α, α, α -trifluoro-2,6-dinitro- N,N -dipropyl-p-toluidine	0.43%
OTHER INGREDIENTS	99.14%
TOTAL 10	00.00%

EPA Reg. No. 10163-346-961 Team® is a Trademark of Gowan Company, LLC EPA Est. 961-PA-01:961-IL-01 Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN **WARNING**

Precautionary Statements Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals.

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Refer to back panel for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.




Distributed by:

Lebanon Seaboard Corporation 1600 E. Cumberland St. • Lebanon, PA 17042 800-233-0628 • (717) 273-1685

NET WEIGHT



Shoes plus socks





Fertilizer with PREEMERGENCE WEED CONTROL

· Goggles, faceshield or safety glasses Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

· Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow people or pets to enter the treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not contaminate water, food or feed by storage or disposal.

Storage and Disposal

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

General Information

General Use Precautions and Restrictions

This product is a granular fertilizer containing Team®, a preemergence herbicide for application to established turfgrass and landscape ornamentals (trees, shrubs, groundcovers, and flowers) for control of annual grasses and broadleaf weeds

Do not apply this product to golf course putting greens.

almond

apricot

ash, white

blackgum

cottonwood

douglas fir

fir, balsam

honeylocust

locust, black

maple, red

oak, pin

peach

pine, red

plum

pine, scotch

sweetgum

sycamore

Annual Grasses

tuliptree

oak, scarlet

larch, Japanese

maple, Norway

cherry

bald cypress

apple, crabapple

birch, European white

chestnut, Chinese

dogwood, flowering

dogwood, kousa

hemlock, Canada

· Do not apply to dichondra. • Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product

Reseeding: Do not reseed treated areas for at least 8 weeks after a single application of 174 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 174 lb of product per acre rate is applied, or 10 to 12 weeks after the last 233 lb of product per acre rate is applied unless injury can be tolerated.

> This product is recommended for use on the following turfgrass species: **Established Warm Season Turfgrass**

Established Cool Season Turfgrass bentarass fairways bahiagrass fescues Bermudagrass Kentucky bluegrass centipedegrass perennial rvegrass fescue, tall St. Augustinegrass

zoysiagrass This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees Common Name Scientific Name

Prunus dulcis Malus spp. Prunus armeniaca Fraxinus americana Taxodium distichum Betula pendula Nyssa sylvatica Prunus spp. Castanea mollissima Populus deltoides Cornus florida Cornus kousa

Pseudotsuga menziesii Abies balsamea Tsuga canadensis Gleditsia triacanthos Larix kaempferi Robinia pseudoacacia Acer platanoides Acer rubrum Acer saccharinum

maple, silver maple, sugar Acer saccharum Quercus palustris Quercus coccinea Prunus persica pine, Austrian Pinus nigra pine, eastern white Pinus strobus pine, Japanese black Pinus thunbergiana pine, loblolly Pinus taeda Pinus resinosa Pinus sylvestris planetree, London Platanus acerifolia Prunus spp. redbud, eastern Cercis canadensis spruce, Colorado Picea pungens spruce, Norway Picea abies spruce, white Picea glauca Liquidambar styraciflua Platanus occidentalis Liriodendron tulipifera walnut, black Juglans nigra

Common Name andromeda, Japanese arborvitae, American azalea barberry, Japanese barberry, mento boxwood, common boxwood, littleleaf camellia, Japanese camellia, sasangua

Ornamental Shrubs

cherrylaurel, American cinquefoil cleyera, Japanese cotoneaster, cranberry cotoneaster, zabel deutzia elaeagnus, silverberry euonymus, spreading euonymus, winged euonymus, wintercreeper firethorn forsythia guava, pineapple

honevsuckle India hawthorn iuniper laurel, mountain lilac, common mockorange pittosporum, Japanese privet redcedar, eastern rhododendron spiraea, vanhoutte viburnum weigela willow vew, angloiap

holly

Pieris japonica Thuja occidentalis Rhododendron spp. Berberis thunbergii Berberis mentorensis Buxus sempervirens Buxus microphylla Camellia japonica Camellia sasangua Prunus caroliniana Potentilla spp. Cleyera japonica Cotoneaster apiculata Cotoneaster zabelii Deutzia spp. Elaeagnus pungens Euonymus kiautschovica Euonymus alatus Euonymus fortunei Pyracantha spp. Forsythia spp. Feijoa sellowiana llex spp. Lonicera spp. Raphiolepis indica Juniperus spp.

Scientific Name

Kalmia latifolia Syringa vulgaris Philadelphus spp. Pittosporum tobira Ligustrum spp. Juniperus virginiana Rhododendron spp. Spiraea vanhouttei Viburnum spp. Weigela spp. Salix spp. Taxus media Taxus cuspidata Podocarpus macrophyllus **Ornamental Groundcover Plantings Common Name** aaronsbeard bellflower, adriatic bellflower, poscharsky ceanothus

coreopsis cotoneaster covote brush crown vetch daisy trailing African fern. asparagus gazania germander ice plant, largeleaf ivy, Algerian ivv. English

lily-of-the-nile lilyturf, bigblue marigold myoporum plumbago, dwarf rockrose rosemary rupturewort snow-in-summer speedwell St. Johnswort stonecrop (sedum)

strawberry, beach thrift verbena wirevine, creepina yarrow, wooly

Scientific Name **Common Name** Hypericum calycinum African daisy aster (perennial) Campanula elatines Campanula poscharskyana balsam Ceanothus spp. blackeved susan

Coreopsis spp. Cotoneaster spp. Baccharis pilularis Coronilla vana Osteospermum fruticosum Asparagus densiflorus Gazania spp. Teucrium chamaedrys Carpobrotus edulis Hedera canariensis Hedera helix Agapanthus spp. Liriope muscari Tagetes spp. Myoporum laetum Ceratostigma umbaginoides Cistus spp.
Rosmarinus officinalis Herniaria glabra

Veronica spp.

Hypericum coris

Sedum spp. Fragaria chiloensis

Muehlenbeckia axillaris

Armeria maritima

Verbena spp.

dusty miller floss flower forget-me-not four o'clock gaillardia gladiolus golden glow impatiens ixora lobelia Cerastium tomentosum lupine

calendula

carnation

coreopsis

cornflower

cosmos

dianthus

dahlia

centaurea, velvet

chrysanthemum

Weeds Suppressed

Weeds suppressed when applied prior to weed germination:

Common Name marigold marigold, cape morningglory nasturtium nicotiana petunia phlox pincushion flower poppy, California portulaca rose salvia shasta daisy snapdragon snow-on-the-mountain stock sunflower sweet alyssum sweet pea sweet sultan sweet william vinca varrow zinnia

Roses and Other Established Flowers

Weeds Controlled

vew. Japanese

vewpine

Weeds controlled when applied prior to weed germination:

Common Name annual bluegrass Poa annua barnyardgrass (watergrass) crabgrass, large or hairy Digitaria sanguinalis crabgrass, smooth Eleusine indica goosegrass (silver crabgrass or crowfoot) green foxtail Setaria viridis yellow foxtail

Scientific Name Echinochloa crus-galli Digitaria ischaemum Setaria glauca

Broadleaf Weeds Common Name chickweed, common knotweed, prostrate lambsquarters, common pigweed purslane, common

Scientific Name Stellaria media Polygonum aviculare Chenopodium album Amaranthus spp. Portulaca oleracea

Common Name creeping spurge creeping woodsorrel panicum, Texas prostrate spurge pusley, Florida spotted spurge woodsorrel

Scientific Name Euphorbia serpens Oxalis corniculata Panicum texanum Euphorbia humistrata Richardia scabra Euphorbia maculata Oxalis spp.

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Uses **Established Turfgrass**

Application Timing

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended period of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

Broadcast Application Rates Cool Season Turfgrass Warm Season Turfgrass Summer annual grass weeds 174 to 349 lb/acre 349 lb/acre (spring application) (4 to 8 lb/1000 sq ft) (8 lb/1000 sq ft) Winter annual grass weeds 174 to 349 lb/acre 349 lb/acre (4 to 8 lb/1000 sq ft) (fall application) (8 lb/1000 sq ft)

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 174 lb plus 174 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 174 lb per acre rate and 10 to 12 weeks following the 233 lb per acre rate. **Specific Use Restrictions**

Do not make more than 2 applications of this product per year on turfgrass.
 Maximum Single Application Rate: 349 lb of Turf Fertilizer – Contains Team Pro 0.86% per acre.
 Minimum Re-Treatment Interval: Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.

• This product must be watered in as soon as possible after application.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

Broadcast Application Rate Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

Specific Use Restrictions

 Do not make more than 2 applications of this product per year on all ornamental plants. Non-bearing fruit and nut trees, non-bearing vineyards:

o Maximum Single Application Rate: 8 lb of this product - Contains Team Pro 0.86% per 1000 sq ft. o Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited

 Landscape ornamentals: o Maximum Single Application Rate: 8 lb of this product - Contains Team Pro 0.86% per 1000 sq ft.

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as

unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed **Limitation of Remedies**

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or 2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.