CONFRONT* 3 & DIMENSION FERTILIZER WITH

LEBANON ProScape®

2,4-D	GROUP	4	HERBICIDE
TRICLOPYR	GROUP	4	HERBICIDE
FLUROXYPYR	GROUP	4	HERBICIDE
DITHIOPYR	GROUP	3	HERBICIDE

LEBANON

ProScape®

FERTILIZER WITH CONFRONT* 3 & DIMENSION

19-0-6

MESA

GUARANTEED ANALYSIS

Total Nitrogen (N) 19%
2.5% Ammoniacal Nitrogen
1.6% Water Insoluble Nitrogen*
12.3% Urea Nitrogen
2.6% Other Water Soluble Nitrogen*
Soluble Potash (K2O) 6%
Sulfur (S) 3.0%
3.0% Combined Sulfur (S)
Derived from: Urea, Methylene Ureas, Ammonium Sulfate, Muriate of Potash.
*4.2% Slowly Available Nitrogen from Methylene Ureas.
Chlorine (Cl) not more than 3%

0 69585 53986 2
FS99 22-53986
B06

SUGGESTED SPREADER SETTINGS

Spreader	LBS/1,000 Sq. Ft. 3.5	4	4.5	5
Andersons SR2000	K 1/2	L	L 1/2	M
Lesco (letter/numeric dial)	R1/4	Q/15	H/16	I/17
Scotts R8	I 1/2	I/17	J/21	K/22
Spyker Ergo-Pro	R1/3	J	J 1/2	K
Chapin SureSpread	12	13	14	16
Earthway Rotary	12	13	14	16

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

If your spreader is not listed, or if you need a spreader setting, please call customer service at **1-800-233-0028**, Monday through Friday, 8 am. To 5 pm.

For the postemergence control of annual and perennial broadleaf weeds, and preemergence control of annual grassy weeds and labeled broadleaf weeds, in established ornamental turf, lawns, golf courses, roadsides and cemeteries.

ACTIVE INGREDIENTS: BY WT.%

Dimethylamine salt of 2,4-D*	0.819%
Triethylamine salt of triclopyr	0.303%
Methylheptyl ester of fluroxypyr	0.177%
Dithiopyr	0.140%
OTHER INGREDIENTS:	98.559%
TOTAL	100.000%

Acid Equivalents:

2,4-dichlorophenoxyacetic acid – 0.680% 13.60 lbs./ton	2,4-D CAS, No. 94-75-7
triclopyr: 3,5,6-trichloro-2-pyridyloxyacetic acid = 0.217% 4.34 lbs./ton	Triclopyr CAS, No. 55335-06-3
fluroxypyr: 4-amino-3,5-dichloro-6-fluoro-2-pyridyloxyacetic acid = 0.127% 2.46 lbs./ton	Fluroxypyr CAS, No. 69377-81-7
dithiopyr – 0.140% 2.8 lbs./ton	Dithiopyr CAS No. 97886-45-8

*Isomer Specific by AOAC Method No. 6.275-6.279 (13th Ed.)

(L) (D)

EPA Reg No. 961-406 **EPA Est. No. 961-PA-01; 961-IL-01**

*Trademark of Corteva Agriscience Superscript used to first letter of lot number.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See back panel for First Aid and additional Precautionary Statements

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.
Before buying this product, read the Warranty Section elsewhere on this label.
If terms are not acceptable, return the unopened container to the place of purchase at once.

In New York State: This product may be applied only by commercial applicators at no more than 241 pounds (0.35 lb. of active ingredient of dithiopyr) per acre (5.5 lbs per 1,000 sq.ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.

LebanonTurf

Manufactured by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
800-233-0628 • (717) 273-1685
www.LebanonTurf.com

NET WEIGHT **40** LBS. (18.14 KG)

COVERS UP TO: 11,400 SQ. FT.

CONFRONT* 3 & DIMENSION FERTILIZER WITH
LEBANON ProScape®

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wear long-sleeved shirt and long pants, socks, shoes, waterproof gloves and protective eyewear (goggles, face shield, or safety glasses). Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE).
All loaders, applicators, and other handlers must wear:
- Long-sleeved shirt and long pants.
- Socks plus shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations
Users should: • Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
• Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **(888)-209-1368** for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.
The product contains chemicals that have properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water of groundwater.
GROUNDWATER ADVISORY: This chemical has properties and characteristics associated with chemicals detected in groundwater. The chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
SURFACE WATER ADVISORY: The product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several weeks after application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
NOT FOR USE ON TURF BEING GROWN FOR SALE OR OTHER COMMERCIAL USE AS SOD; OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES.

GENERAL INFORMATION

Lebanon ProScape® Fertilizer with Confront* 3 & Dimension provides broad-spectrum postemergence control of actively growing annual and perennial broadleaf weeds, preemergence or early postemergence control of certain annual grassy weeds, and preemergence control or suppression of certain broadleaf weeds in established turf, including residential lawns, golf courses, roadsides and cemeteries. It is also labeled for use in sod farms and other commercial and residential turf areas. Apply when broadleaf weeds are actively growing and prior to germination or tillering of target annual grassy weeds. Application under drought conditions may provide less than desirable results. Applications to moist foliage helps granules adhere to weed surface and will improve postemergence control.
Control of Crabgrass Preemergence and Early Postemergence Control
Preemergence and postemergent control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass. In addition, it will control or suppress annual bluegrass, barnyardgrass, cinnefoilgrass, foxtail (green and yellow), goosegrass, kikuyugrass, ryegrass (annual and perennial) and smartweed.
This product also provides early postemergent control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergent crabgrass control is a 2-4 week longer period of time (window of application) during which crabgrass control can be achieved.

WEED RESISTANCE MANAGEMENT

For resistance management, **Lebanon ProScape® Fertilizer with Confront* 3 & Dimension** is a Group 3 and Group 4 herbicide. Any weed population may contain or develop certain plants naturally resistant to **Lebanon ProScape® Fertilizer with Confront* 3 & Dimension** and other Group 3 and/or Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies listed below should be followed.
To delay herbicide resistance, take one or more of the following steps:
Do not apply **Lebanon ProScape® Fertilizer with Confront* 3 & Dimension** or other Group 3 and Group 4 herbicides within a growing season with different herbicide groups that control the same weeds.
Adopt an integrated weed management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers outbreak, biological varieties and other turf management practices.
Scout after herbicide application to monitor weed populations for early signs of resistance development. Indications of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species.
If resistance is suspected, prevent weed seed production in the affected area by using an alternative herbicide from a different group or by a mechanical weed removal method. Prevent movement of resistant weed seeds to other areas by cleaning equipment thoroughly.
Rotate the use of **Lebanon ProScape® Fertilizer with Confront* 3 & Dimension** or other Group 3 and Group 4 herbicides with different herbicides that control the same weeds.
Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance management and/or integrated weed management recommendations for specific types of plants and weed biotypes.

PRECAUTIONS

Be sure that use of this product conforms to all applicable regulations. **Avoid off target application.** Applications should be made so as to avoid spreading material off target. Avoid getting granules in shrub bed or other areas where they may seriously injure susceptible plants.

RESTRICTIONS

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
DO NOT use this product where desirable clovers are present or apply (spread) the granules onto flowers, vegetables or ornamental shrubbery.
DO NOT apply this product through any type of irrigation system.
DO NOT exceed specified dosage for any area, be particularly careful within dripline of trees and other ornamental species.
DO NOT use on grass species such as bentgrass or St. Augustinegrass unless injury can be tolerated. Use of this product on certain varieties of fine fescue may result in undesirable turf injury.
DO NOT apply to newly seeded areas until after the first spring of the first year of establishment until the turfgrass is fully green and has established new roots.
DO NOT use **LEBANON PROSCAPE FERTILIZER WITH CONFRONT* 3 & DIMENSION** on golf course putting greens or tees.
DO NOT contaminate domestic or irrigation waters.
DO NOT apply on ditches used to transport irrigation water.
DO NOT apply under conditions that may have a potential to run off to surface water or adjacent land. **DO NOT** apply where runoff or irrigation water may flow onto susceptible crops as injury may result.
Where possible, use methods that reduce soil erosion, such as strip seeding; these methods also reduce pesticide run-off. Use of vegetative filter strips is recommended along rivers, creeks, streams, wetlands, etc.
DO NOT use clippings from treated areas for mulching vegetables or fruit trees.

- This product is persistent and may be present in treated plant materials for over 30 days after application. **DO NOT** remove grass clippings off-site for compost distribution or mulching until 30 days after application.
- **DO NOT** apply more than 5.0 lbs. of this product (0.071 lb. triclopyr a.e. and 0.0072 lb. dithiopyr) per 1,000 sq. ft. per application i.e., 5.47 lb. triclopyr a.e. and 0.315 lb. dithiopyr per acre.
- **DO NOT** apply more than 10.0 lbs. of this product (0.071 lb. triclopyr a.e. and 0.014 lb. dithiopyr) per 1,000 sq. ft. per year, i.e., 0.946 lb. triclopyr a.e. and 0.632 lb. dithiopyr per year.
- Minimum retreatment interval is 28 days.

Maximum Use Rate Restrictions in Sod Belt States: This product may be applied only by commercial applicators at no more than 241 pounds (0.35 lb. of active ingredient of dithiopyr) per acre (5.5 lbs per 1,000 sq. ft.) per year. Use of this product in Nassau and Suffolk Counties is prohibited.
Re-entry Restrictions: DO NOT allow people other than applicator(s) or pets or treatment area during application. **DO NOT** enter treatment areas until dust has settled.
Seeding Restrictions: DO NOT reseed for 10 weeks after application.
Reinstatement Restrictions: DO NOT apply more than 2 broadcast applications per year per treatment site. To minimize the potential for grass injury, additional applications should not be made within 5 weeks of a previous application.

LEBANON PROSCAPE FERTILIZER WITH CONFRONT* 3 & DIMENSION is recommended for use on the following established turfgrass species:

Established Cool Season Turf Grasses		Established Warm Season Turf Grasses*	
Common Name	**Scientific Name**	**Common Name**	**Scientific Name**
bentgrass*	*Agrostis Species*	Bahiagrass	*Paspalum notatum var saurae parody*
bluegrass, Kentucky	*Poa pratensis*	Bermudagrass	*Cynodon dactylon*
bluegrass, rough	*Poa trivialis var. communiata*	Centipedegrass	*Eremochloa ophiuroides*
fescue, creeping red	*Festuca rubra*	Centipedegrass	*Eremochloa ophiuroides*
fescue, sheeps	*Festuca rubra*	Zoysiagrass	*Zoysia japonica*
fescue, tall	*Festuca arundinaceae*	Zoysiagrass	*Zoysia matrella*
ryegrass, perennial	*Lolium perenne*	St. Agustinegrass (except Floratam')	*Stenotaphrum secundatum*
		fescue, tall (growing in warm season areas)	*Festuca arunicaceae*

*Use **LEBANON PROSCAPE FERTILIZER WITH CONFRONT* 3 & DIMENSION** on this species only if some injury can be tolerated.

WHEN TO APPLY

Established Lawns: Apply when broadleaf weeds are actively growing and prior to germination or tillering of annual grasses. **LEBANON PROSCAPE FERTILIZER WITH CONFRONT* 3 & DIMENSION** may be applied as a single application, as a split application or as a sequential application in the spring, summer or fall as described in the following use programs. Use of split applications 8-10 weeks apart may provide improved weed control.
Newly Seeded Areas: When establishing a new lawn of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied 2 weeks after the first sign of grass germination. For other turfgrass varieties, wait until the grass is well-established (3 mowings) before applying this product.

HOW TO APPLY

For optimum broadleaf weed control, apply to recently mowed turf when foliage is moist. When possible, delay mowing or irrigation for several days after treatment. The dithiopyr preemergent component in this product is not effective until activated by rainfall or irrigation. Performance is improved by 1/2 inch of rainfall or sprinkler irrigation within a 30 day period. Erratic weed control may result if not activated by rainfall or irrigation.
Uniform coverage is essential for best results. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Avoid streaking, skips or overlaps. Adjust the spreader on the spreader delivers the appropriate application rate as recommended below.

APPLICATION RATES

Apply 152 to 218 lbs. per acre (3.5 to 5 lbs. per 1,000 sq. ft.) of **LEBANON PROSCAPE FERTILIZER WITH CONFRONT* 3 & DIMENSION** to control actively growing broadcleaf weeds and prevent germination of labeled annual grasses and broadleaf weeds growing in established turf. Broadleaf and grassy weed species germinate at different times. Only emerged weeds present at time of application are controlled.
Specific rate recommendations are based on the user's location, the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the users' location within a region. The following charts show application rates and some of the common broadleaf weeds controlled. The rates vary based on region which are described as follows:

Transition: DE, KS, KY, MD, MO, NJ, VA, southwestern PA, southeastern areas of IL, IN, OH, and coastal areas of CT & RI
Cool: New England, northern PA, and northern coastal areas of IL, IN, LA, MI, NC, SC, and TN
Coastal: Southern areas of IL, and southern coastal areas of IL, IN, LA, MS, NC, SC, and TX

POSTEMERGENCE & PREEMERGENCE PROGRAMS WITH APPLICATION RATES & WEEDS CONTROLLED

152 LBS./ACRE;	Fluroxypyr AI/Acre	Triclopyr AI/Acre	2,4-D AI/Acre	Dithiopyr A.I./Acre
3.5 LBS./1,000 SQ. FT.	0.147	0.330	1.054	0.0294

Preemergent control of the following broadleaf weeds and 2.5 months preemergent control, as well as early postemergence control of crabgrass up to the 3 leaf or first tiller stage, at sites with high soil turf* or all states except MA, parts of states not included in the transition, south, coastal south, or west.

bedstraw	chamise	cornflower	field bindweed	lespedeza	shepherdspurse	waterbell
bittercress	chickweeds	creeping beggarweed	geranium, carolina	little starwort	spanbeed	witch
black medic	chicory	dandelion	goldenrod	mallow	smartweed	woodsorrel
bottlebrush, virginia	cinquefoil	daylilyear	henbit	matchweed	snakeroot	yarrow
cocklebur	clovers	deadnettle, purple	hop clover	pineappleweed	spiderwort	
corn speedwell	cocklebur	dock	knotweed	plantain	spotted spurge	
cudweed	coffeeweed	dwarf beggarweed	lambsquarters	purslane	starchweed	

174 LBS./ACRE;	Fluroxypyr AI/Acre	Triclopyr AI/Acre	2,4-D AI/Acre	Dithiopyr A.I./Acre
4.0 LBS./1,000 SQ. FT.	0.214	0.378	1.183	0.252

Preemergent control to include the above broadleaf weeds listed below and provides 4-5 months of preemergent control on lawns* turf** where weed control has been conducted during the previous year. Also provides early postemergent control of crabgrass up to the 3 leaf or first tiller stage on high cut turf**. All states and parts of states not included in the south, coastal south, or west. Recommend split applications of 4.0 + 4.0 lbs. on high cut turf** in the coastal south and lowcut turf in the south.

burdock	compositeae<	knotweed	oxalis (citricila and corniculata)<	poison ivy	parsley-piert<	wild carrot
buttercup	ground ivy<	mustard	plantain	poison oak	wild violet<	veronica species< (suppression)

*Discontinued 2 applications, 2 to 6 weeks apart, apply to moist foliage.

196 LBS./ACRE;	Fluroxypyr AI/Acre	Triclopyr AI/Acre	2,4-D AI/Acre	Dithiopyr A.I./Acre
4.5 LBS./1,000 SQ. FT.	0.241	0.426	1.332	0.284

Provides postemergent control and provides 5.5 months of preemergent control on lawns* turf** where weed control has been conducted during the previous year. Provides early postemergent control of crabgrass up to the 3 leaf or first tiller stage on high cut turf**. Includes the West use higher rate in local areas with longer crabgrass seasons, see below. Note precautions for warm season turf species.

218 LBS./ACRE;	Fluroxypyr AI/Acre	Triclopyr AI/Acre	2,4-D AI/Acre	Dithiopyr A.I./Acre
5.0 LBS./1,000 SQ. FT.	0.268	0.473	1.482	0.316

Improved postemergent control and provides 4-6 months of preemergent control on lawns* turf** where weed control has been conducted during the previous year in the coastal south and west. For coastal south split applications of 5.0 + 5.0 lbs. are recommended. Note precautions for warm season turf species.

*high cut turf such as home lawns and commercial sites.
**Lowcut turf such as golf fairways.

STORAGE AND DISPOSAL

DO NOT contaminate water, feed, or feed by storage or disposal.
PESTICIDE STORAGE: Store product in its original container in a cool, dry, secure area out of the reach of children and animals
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at any approved waste disposal facility.
CONTAINER DISPOSAL: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.
By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.
CONDITIONS: The directions for use of this product are based on tests believed to be reliable and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Lebanon Seaboard Corporation. All such risks shall be assumed by the user or buyer.
DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, LEBANON SEABOARD CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Lebanon Seaboard Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Lebanon Seaboard Corporation DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.
LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LEBANON SEABOARD CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

For technical assistance or more information about our products, visit LebanonTurf.com
Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-4ltexas.com