

LEBANON ProScape®

FERTILIZER

WITH **CONFRONT®** HERBICIDE
BROADLEAF WEED CONTROL

19-0-7

GUARANTEED ANALYSIS

Total Nitrogen (N) **19.0%**
 2.5% Ammoniacal Nitrogen
 1.6% Water Insoluble Nitrogen*
 12.3% Urea Nitrogen
 2.6% Other Water Soluble Nitrogen*

Soluble Potash (K₂O) **7.0%**
 Sulfur (S) 3.0%
 3.0% Combined Sulfur (S)
 Iron (Fe) 1.0%
 0.10% Water Soluble Iron (Fe)

Derived From: Urea, Methylene Ureas, Ammonium Sulfate, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.

*4.2% Slowly available Nitrogen from Methylene Ureas.
 Chlorine (Cl) not more than 5.0%



Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.html>

SUGGESTED SPREADER SETTINGS					
Spreader	135.5#/acre	156.4#/acre	Spreader	135.5#/acre	156.4#/acre
LebanonTurf	4½	5	Lesco (letter/Numeric)	I/17	J/21
Andersons AccuPro	K	M	PennMulch HVO	L	N
Earthway Rotary	15	17	ProScape SS	L	N
Gandy	27	29	Spyker	4½	5
Lely	5½ II	6 II	Vicon (all models)	25	28

MESA® is a registered trademark of Lebanon Seaboard Corporation.

To be applied only under the direct supervision of licensed pesticide applicators responsible for turf weed control programs.

- Selectively controls annual and perennial broadleaf weeds
- Provides a feeding of fertilizer

ACTIVE INGREDIENT:

triclopyr: 3,5,6 trichloro-2-pyridinyloxyacetic acid, triethylamine salt 0.50%
 clopyralid: 3,6-dichloro-2-pyridinyloxyacetic acid, triethylamine salt 0.18%
OTHER INGREDIENTS: 99.32%
TOTAL 100.00%

Acid equivalent: (L) (D)
 triclopyr - 0.36% EPA Est. 961-PA-01; 961-IL-01
 clopyralid - 0.12% Superscript used is first letter of lot number.
 EPA Reg. No. 62719-262-961
®Trademarks of Corteva Agriscience and its affiliated companies

**KEEP OUT OF REACH OF CHILDREN.
WARNING**

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

**Precautionary Statements
Hazards to Humans and Domestic Animals**

WARNING

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through The Skin

Do not get in eyes or on clothing. Avoid contact with skin. Wear long-sleeved shirt and long pants, socks and shoes, protective eyewear and chemical resistant gloves made of rubber or other waterproof material. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

User Safety Recommendations

- Users should:
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.



Manufactured by:
Lebanon Seaboard Corporation
 1600 E. Cumberland St. • Lebanon, PA 17042
 800-233-0628 • (717) 273-1685
www.LebanonTurf.com

NET WEIGHT **40** LBS. (18.14 Kg)

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractured surfaces, and substrates which could allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in a clean, dry area away from extremes of heat and cold and out of direct sunlight.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Product Information

This product is a broad spectrum weed killer that will control broadleaf weeds in established cool season and warm season turfgrass.

This product may be used on the following turf species:

Common Name	Established Cool Season Turfgrass	Scientific Name
bluegrass, Kentucky		Poa pratensis
fescue, chewing		Festuca rubra var.commutata
fescue, creeping red		Festuca rubra
fescues, sheeps		Festuca ovina
fescue, tall		Festuca arundinaceae
ryegrass, perennial		Lolium perenne
	Established Warm Season Turfgrass¹	
bahiagrass (Argentine or Pensacola)		Paspalum spp.
buffalograss		Buchloe dactyloides
centipedegrass		Eremochloa ophiuroides
bermudagrass ²		Cynodon dactylon
fescue, tall (growing in warm seasonarea)		Festuca arundinaceae
zoysiagrass		Zoysia japonica
zoysiagrass		Zoysia tenuifolia
	¹ Do not treat warm season turfgrasses with Turf Fertilizer Contains Confront when mowing height is less than 1/2 inch.	
	² Do not apply Confront to bermudagrass on sod farms.	

Use Precautions:

This product may discolor and/or stunt turf that is not well-established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply this product only to healthy, well-established turf that has a well-anchored root system. Bermudagrass, centipedegrass, zoysiagrass, and bahiagrass may be temporarily discolored by this product.

Use Restrictions

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

In **California, Oregon and Washington**, turfgrass and lawn uses are restricted to golf courses only.

Do not use on residential turf. Turfgrass and lawn uses are restricted to non-residential sites.

Apply this product only as specified on this label.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled.

Do not apply to exposed roots of shallow rooted trees and shrubs and allow to contact exposed suckers or injury may occur.

Do not apply this product directly to, or otherwise permit it to come into direct contact with, cotton, grapes, tobacco, vegetables, flowers, fruit or ornamental trees, or other nontarget broadleaf plants.

Do not reseed until 4 weeks after application.

Do not use this product on golf course putting greens and tees.

Do not send grass clippings to a compost facility.

Do not collect grass clippings for mulch or compost.

Applicator must give notice to landowners/property managers to not use grass clippings for composting.

Do not apply on ditches used to transport irrigation water.

Do not contaminate water used for irrigation or domestic purposes.

Do not apply where runoff or irrigation water may flow onto susceptible crops as injury may result.

Do not contaminate feed or foodstuffs.

Do not graze treated areas. Do not feed clippings to livestock.

Do not apply when weather conditions favor drift from target areas. Drift or runoff may adversely affect nontarget plants.

Do not apply at greater than specified rates, especially under the branch spread (dripline) of trees.

Do not seed, sod, or sprig after application. Delay until no sooner than 4 weeks after application.

Do not apply this product to bentgrass; hybrid bermudagrass varieties such as tifway, tifdwarf, and tifgreen; St. Augustinegrass; and ground covers such as lippia.

On bahiagrass and centipedegrass, do not apply before mid-March or until after the second mowing of the year.

In the fall, do not treat centipedegrass after the first of October or bahiagrass after the first November.

• Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>.

• Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf.

Treatment of Turfgrass Species Not Listed on the this product Label

Users who wish to use this product on turfgrass species not listed on this label may determine the suitability for such uses by treating a small area at the specified rate. Prior to treatment of larger areas, observe the treated area for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the treatment is safe to the target species. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not listed on this label.

Application Directions

Use equipment that will ensure uniform coverage and apply 3.58 lbs per 1000 sq feet (156.46 lb per acre, equivalent to 0.187 lb ae/acre clopyralid and 0.56 lb ae/acre of triclopyr) and 0.68 lbs of Nitrogen per 1000 sq ft when weeds are actively growing. Good soil moisture conditions should be present before application to achieve optimal results. Application under drought conditions may provide less than desirable results. Broadleaf weed species germinate at different times. Only emerged weeds present at time of application are controlled. To minimize the potential for turfgrass injury, do not make additional applications within 4 weeks of a previous application. Apply to newly seeded or sprigged turfgrass only after the fourth mowing or after the turfgrass seedlings have tillered and developed an adequate secondary root system.

- Do not apply more than 2 applications per year and do not use more than 0.375 lb ae clopyralid/1.125 lb ae triclopyr (312.5 lbs of this product) per acre per year.
- In Florida and New York, the maximum use rate is 0.25 lb ae clopyralid/0.75 lb ae triclopyr per acre (205.6 lbs of this product) per acre per year.

Weeds Controlled

Apply this product to moist or dry foliage at 3.58 lbs per 1000 sq feet (156.46 lb per acre, equivalent to 0.187 lb ae/acre clopyralid and 0.56 lb ae/acre of triclopyr) to control the following weeds:

black medic	hawkweed	burdock	lespedeza
bristly hawksbeard	hop clover	Canada thistle	matchweed
broadleaf plantain	narrow leaf plantain (buckhorn)	coffee weed	mouseear chickweed
catsear (coast dandelion)	red clover	common chickweed	musk thistle
common dandelion	white clover	common cocklebur	old world diamond flower
false dandelion		common ragweed	poison ivy
		common vetch	prostrate spurge
		common yellow woodsorrel	roundleaf mallow
		creeping beggarweed	sheep sorrel
		creeping woodsorrel	sheepspurse
		curly dock	slender speedwell
		dollarweed	smartweed
		dwarf beggarweed	spurweed
		English lawn daisy	Virginia buttonweed
		goldenrod	Virginia pepperweed
		ground ivy	wild buckwheat
		henbit	wild violet
		lambquarters	

Note: Moist foliage application will provide quicker results. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present.

For harder to control weeds, a repeat application in 4 to 6 weeks may be necessary.

Weed Resistance Management:

Triclopyr and clopyralid, the active ingredients in this product, are Group 4 herbicides based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plants resistant to Group 4 herbicides. Resistant weeds may dominate the weed population if these herbicides are used repeatedly in the same field. Such resistant weed plants may not be effectively managed using Group 4 herbicides but may be effectively managed utilizing other herbicides alone or in mixtures from a different herbicide Groups that are labeled for control of these weeds and/or by using cultural or mechanical practices. However, a herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds or to report herbicide failures.

Best Management Practices:

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. It is recommended to scout for weeds before Turf Fertilizer Contains Confront application for identification and growth stage and after application to facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all i such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.