

satisfactory control or suppression

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may contact 1-800-992-5994 for emergency medical treatment information.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye

All mixers, loaders, applicators and other handlers must wear:

Personal Protective Equipment (PPE)

	Rate	
pressed	lb/acre	lb/1000 sq ft
ed, chicory, clover, dandelion, dock, dollarweed, eclipta, healall ,henbit, ivy,(ground), nedic (black), morningglory, peppergrass, pigweed, plantain, plantain (broadleaf), ne, ragweed, sheepsorrel, shepherd's purse, speedwell, thistle	150-175	3.4 - 4

This product, which contains the active ingredient penoxsulam is a Group 2 herbicide based on the mode of action classification system of the Weed

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program

To aid in the prevention of developing weeds resistant to this product, users should:

Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.

If using post-emergence herbicides or tank mixes, control weeds early when they are relatively small.

Apply full rates of this product for the most difficult to control weed in the field at the specified time to minimize weed escapes

Scout fields after application to detect weed escapes or shifts in control of weed species. Control weed escapes before they reproduce by seed or proliferate vegetatively.

Report any incidence of non-performance of this product against a particular weed to your local company representative, local retailer, or county extension agent. Contact your local company representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective modes of action for each target weed

If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 2 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.

Suspected herbicide-resistant weeds may be identified by these indicators:

§ Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; § A spreading patch of non-controlled plants of a particular weed species; and

Surviving plants mixed with controlled individuals of the same species

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

Use a broad spectrum herbicide with other mode of action as a foundation in a weed control program, if appropriate

Avoid making more than two sequential applications of this product and any other Group 2 herbicides within a single growing season unless mixed with an herbicide with a different mode of action with an overlapping spectrum for the difficult-to-control weeds.

Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an

Use good agronomic principles that enhance crop development and crop competitiveness.

Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

To optimize product performance, mow the turfgrass two to three days before applying this product. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of at least 1/2 inch may be needed at least one to two days before application. For optimal results, apply this product in the morning when dew is present. If necessary, sprinkler irrigate with less than 1/4 inch of water just prior to application to maximize particle adherence to leaf surfaces. Avoid watering of reated area for 24 hours following application. Avoid making applications if rainfall is expected within 24 hours after treatmen

Application Techniques for Applying Lebanon ProScape Fertilizer with LockUp® Herbicide

When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution

• A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent

• For small, difficult to treat areas use a shaker-type applicator made from a small container with holes punched in the bottom. Carefully measure the amount of

. For treating smaller areas, use a hand held or push-type rotary applicator such as a whirlybird or cyclone unit. For hand held units, walk and turn the crank at a

Rates and Timing of Application

Apply 150 to 175 lb of this product per acre (3.4 to 4 lb per 1000 sq ft) per application to established turfgrass. Possible re-treatment in approximately 30 days may be necessary if turfgrass is heavily infested with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed 350 lb of this product per acre (8 lb per 1000 sq ft) per year because damage to turfgrass may occur. Do not make more than two applications per season

Precaution: When applying to warm season turfgrass under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. Reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of turfgrass.

Best weed control results are obtained when this product is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times. This product may also be applied to newly established warm season turfgrass. Delay treatments four weeks after sodding, sprigging or plugging.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer,

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by buyer. Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

(1)Refund of purchase price paid by buyer or user for product bought, or (2) Replacement of amount of product used

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses

The terms of the Warranty Disclaimer. Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of

Produced by: Lebanon Seaboard Corporation 1600 E Cumberland St • Lebanon, PA 17042

rev: BC a/24

Net Weight 40 LBS. (18.14 kg)