

LebanonTurf

Country Club®

ROOT REVIVER

Quick aerification recovery
Speeds up root establishment
Increases turf density

GRANULAR FERTILIZER + mycorrhizal fungi 3 - 3 - 4

GUARANTEED ANALYSIS

Total Nitrogen (N)..... 3.0%
1.50% Ammoniacal Nitrogen
1.50% Water Insoluble Nitrogen*
Available Phosphate (P₂O₅) 3.0%
Soluble Potash (K₂O)..... 4.0%
Sulfur (S) 3.0%
3.0% Combined Sulfur (S)
Iron (Fe)..... 1.0%
0.01% Water Soluble Iron (Fe)

Derived From: Wheat Middlings, Poultry Compost, Feather Meal, Diammonium Phosphate, Potassium Sulfate, Ammonium Sulfate, Kelp Meal, Ferrous Sulfate, and Ferric Oxide.

*1.5% slowly available Nitrogen from Poultry Compost and Feathermeal.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS:

0.50% Humic Acids
0.15% Ascorbic Acid (Vitamin C)
0.05% Amino Acid (Glycine)
0.02% Myo-Inositol
0.01% Thiamine (Vitamin B₁)

Mycorrhizal fungi (#viable propagules/gram)

Rhizophagus irregularis 1.2
Rhizophagus clarus 1.2
Rhizophagus aggregatus 0.6
Rhizophagus diaphanus 1.2
Funneliformis mosseae 1.2
Funneliformis monosporus 0.6
Septoglomus deserticola 1.2
Claroideoglonus etunicatum 1.2
Claroideoglonus claroideum 1.2



F699

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)..... 3.0%
1.50% Ammoniacal Nitrogen
1.50% Water Insoluble Nitrogen*
Available Phosphate (P₂O₅) 3.0%
Soluble Potash (K₂O)..... 4.0%
Sulfur (S) 3.0%
3.0% Combined Sulfur (S)
Iron (Fe) (actual) 1.0%
0.01% Water Soluble Iron (Fe)

Derived From: Wheat Middlings, Poultry Compost, Feather Meal, Diammonium Phosphate, Potassium Sulfate, Ammonium Sulfate, Kelp Meal, Ferrous Sulfate, and Ferric Oxide.

*1.5% slowly available Nitrogen from Poultry Compost and Feathermeal.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS:

0.50% Humic Acids
0.15% Ascorbic Acid (Vitamin C)
0.05% Amino Acid (Glycine)
0.02% Myo-Inositol
0.01% Thiamine (Vitamin B₁)

Mycorrhizal fungi (#viable propagules/gram)

Rhizophagus irregularis 1.2
Rhizophagus clarus 1.2
Rhizophagus aggregatus 0.6
Rhizophagus diaphanus 1.2
Funneliformis mosseae 1.2
Funneliformis monosporus 0.6
Septoglomus deserticola 1.2
Claroideoglonus etunicatum 1.2
Claroideoglonus claroideum 1.2



NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 lb. Nitrogen per 1,000 sq. ft.

Spreader	8#/1,000 sq. ft. or 3.9 kg/100m ² (covers up to 5,000 ft ² of lawn)	10#/1,000 sq. ft. or 4.8 kg/100m ² (covers up to 4,544m ² of lawn)	Spreader	8#/1,000 sq. ft. or 3.9 kg/100m ² (covers up to 5,000 ft ² of lawn)	10#/1,000 sq. ft. or 4.8 kg/100m ² (covers up to 4,544m ² of lawn)
LebanonTurf	6	6.5	Lesco (letter/Numeric)	L/24	M/26
Andersons AccuPro	M	O	PennMulch HVO	P	Q
Earthway Rotary	20	24	ProScape SS	P	Q
Gandy	32	36	Spyker	6	6.5
Lely	6.5 ll	7 ll	Vicon (all models)	34	40

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Lebanon Seaboard Corporation
1600 E. Cumberland St • Lebanon, PA 17042
800-233-0628 • (717) 273-1685

50 Lbs. (22.7 kg) Net Weight



27-37101

G/23

LebanonTurf

Country Club®

ROOT REVIVER

TURF APPLICATIONS

Aerification:	Apply 8 lbs. per 1,000 sq. ft. (4kg per 100m ²)
Sod / Seed:	Apply 10 lbs. per 1,000 sq. ft. (5kg per 100m ²) In Florida - Apply 8 lbs. per 1,000 sq. ft. Maximum of 2 applications per year)
Divot Mix:	Add 25 lbs. per cubic yard of divot mix

PLANTINGS

Mix thoroughly with the soil being used to fill the hole where being planted.

Trees:	½ lb. (1 cup or 227g) for every 1" diameter of trunk
Shrubs:	One gallon (3.7 liters) containers – ¼ lb. (1/2 cup or 110g) Five gallon (18.9 liters) containers – 1 lb. (2 1/4 cups or 450g)

When used as directed. Country Club Root Reviver will supplement your regular fertilizer program by providing up to .30 lbs. of nitrogen per 1,000 sq. ft. (In Florida - .24 lbs. of nitrogen per 1,000 sq. ft.)

WARNING



Hazard Statements:

May cause skin irritation.
May be harmful if inhaled.
May cause eye irritation.

Precautionary Statements for handling:

Wash hands and exposed skin thoroughly after handling.
Wear protective gloves, protective clothing, eye protection, and face protection.
If eye exposure occurs, rinse with water and seek medical attention.
Take off contaminated clothing and wash before reuse.
If skin irritation occurs seek medical advice
Store locked up in a well-ventilated place.
Keep the container tightly closed and out of reach of children.
Dispose of in accordance with all federal, state and local regulations.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when
1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application
2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State
3) Re-establishing or repairing a turf area.
*In Maryland, the soil testing laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

In Pennsylvania: Do not exceed 0.9 lbs of total nitrogen per 1,000 sq. ft. per application.

In the state of Florida apply fertilizer at rates as described by Best Management Practices for the Enhancement of Environmental Quality on Florida Golf Courses.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Information regarding the content and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>.

Not for use on Food Crops.